

General

Guideline Title

2011 ACCF/AHA guideline for the diagnosis and treatment of hypertrophic cardiomyopathy: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines.

Bibliographic Source(s)

Gersh BJ, Maron BJ, Bonow RO, Dearani JA, Fifer MA, Link MS, Naidu SS, Nishimura RA, Ommen SR, Rakowski H, Seidman CE, Towbin JA, Udelson JE, Yancy CW. 2011 ACCF/AHA guideline for the diagnosis and treatment of hypertrophic cardiomyopathy: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol. 2011 Dec 13;58(25):e212-60. [453 references] [PubMed](#)

Guideline Status

This is the current release of the guideline.

Recommendations

Major Recommendations

Definitions for the weight of the evidence (A-C) and classes of recommendations (I-III) are provided at the end of the "Major Recommendations" field.

Diagnosis

Genetic Testing Strategies/Family Screening

Class I

1. Evaluation of familial inheritance and genetic counseling is recommended as part of the assessment of patients with hypertrophic cardiomyopathy (HCM) (Ho et al., 2002; Arad et al., 2005; Morita et al., 2008; Niimura et al., 1998; Van Driest et al., 2002; Van Driest et al., "Comprehensive," 2004). (Level of Evidence: B)
2. Patients who undergo genetic testing should also undergo counseling by someone knowledgeable in the genetics of cardiovascular disease so that results and their clinical significance can be appropriately reviewed with the patient (Christiaans et al., "Genetic," 2009; Michie et al., 1997; Michie et al., 1998; Offert et al., 2004; Christiaans et al., "Quality," 2009). (Level of Evidence: B)
3. Screening (clinical, with or without genetic testing) is recommended in first-degree relatives of patients with HCM (Ho et al., 2002; Arad et al., 2005; Morita et al., 2008; Niimura et al., 1998; Van Driest et al., "Comprehensive," 2004; Fokstuen et al., 2008; Olivotto et al., 2008). (Level of Evidence: B)
4. Genetic testing for HCM and other genetic causes of unexplained cardiac hypertrophy is recommended in patients with an atypical clinical presentation of HCM or when another genetic condition is suspected to be the cause (Maron et al., 2001; Rosenzweig et al., 1991; Spada et al., 2006). (Level of Evidence: B)

Class IIa

1. Genetic testing is reasonable in the index patient to facilitate the identification of first-degree family members at risk for developing HCM (Ho et al., 2002; Van Driest et al., 2002; Fokstuen et al., 2008). (Level of Evidence: B)

Class IIb

1. The usefulness of genetic testing in the assessment of risk of sudden cardiac death (SCD) in HCM is uncertain (Moolman et al., 1997; Woo et al., 2003). (Level of Evidence: B)

Class III: NO BENEFIT

1. Genetic testing is not indicated in relatives when the index patient does not have a definite pathogenic mutation (Ho et al., 2002; Arad et al., 2005; Morita et al., 2008; Niimura et al., 1998; Van Driest et al., 2002; Van Driest et al., "Comprehensive," 2004; Ho et al., 2000). (Level of Evidence: B)
2. Ongoing clinical screening is not indicated in genotype-negative relatives in families with HCM (Ho et al., 2000; Ingles et al., 2005; Van Driest et al., "Myosin," 2004; Jeschke et al., 1998). (Level of Evidence: B)

Genotype-Positive/Phenotype-Negative Patients

Class I

1. In individuals with pathogenic mutations who do not express the HCM phenotype, it is recommended to perform serial electrocardiography (ECG), transthoracic echocardiography (TTE), and clinical assessment at periodic intervals (12 to 18 months in children and adolescents and about every 5 years in adults), based on the patient's age and change in clinical status (Christiaans et al., "Ventricular," 2009; Andersen et al., 2009; Christiaans et al., 2010; Michels et al., 2009). (Level of Evidence: B)

Electrocardiography

Class I

1. A 12-lead ECG is recommended in the initial evaluation of patients with HCM. (Level of Evidence: C)
2. Twenty-four hour ambulatory (Holter) electrocardiographic monitoring is recommended in the initial evaluation of patients with HCM to detect ventricular tachycardia (VT) and identify patients who may be candidates for implantable cardioverter-defibrillator (ICD) therapy (Maron BJ et al., "American," 2003; Elliott et al., 2006; Maron et al., 1981; Monserrat et al., 2003). (Level of Evidence: B)
3. Twenty-four hour ambulatory (Holter) electrocardiographic monitoring or event recording is recommended in patients with HCM who develop palpitations or lightheadedness (Maron BJ et al., "American," 2003; Elliott et al., 2006; Maron et al., 1981). (Level of Evidence: B)
4. A repeat ECG is recommended for patients with HCM when there is worsening of symptoms. (Level of Evidence: C)
5. A 12-lead ECG is recommended every 12 to 18 months as a component of the screening algorithm for adolescent first-degree relatives of patients with HCM who have no evidence of hypertrophy on echocardiography. (Level of Evidence: C)
6. A 12-lead ECG is recommended as a component of the screening algorithm for first-degree relatives of patients with HCM. (Level of Evidence: C)

Class IIa

1. Twenty-four-hour ambulatory (Holter) electrocardiographic monitoring, repeated every 1 to 2 years, is reasonable in patients with HCM who have no previous evidence of VT to identify patients who may be candidates for ICD therapy (Monserrat et al., 2003). (Level of Evidence: C)
2. Annual 12-lead ECGs are reasonable in patients with known HCM who are clinically stable to evaluate for asymptomatic changes in conduction or rhythm (i.e., atrial fibrillation [AF]). (Level of Evidence: C)

Class IIb

1. Twenty-four-hour ambulatory (Holter) electrocardiographic monitoring might be considered in adults with HCM to assess for asymptomatic paroxysmal AF/atrial flutter. (Level of Evidence: C)

Imaging

Echocardiography

Class I

1. A TTE is recommended in the initial evaluation of all patients with suspected HCM (Maron, 2002; Klues, Schiffers, & Maron, 1995; Wigle et al., 1985; Wigle et al., 1995; Adabag, Kuskowski, & Maron, 2006; Afonso, et al., 2008; Fifer & Vlahakes, 2008; Soor et al., 2009). (Level of Evidence: B)
2. A TTE is recommended as a component of the screening algorithm for family members of patients with HCM unless the family member is genotype negative in a family with known definite mutations (Bos, Towbin, & Ackerman, 2009; Maron, Seidman, & Seidman, 2004; Binder et al., 2006; Hershberger et al., 2009). (Level of Evidence: B)
3. Periodic (12 to 18 months) TTE screening is recommended for children of patients with HCM, starting by age 12 years or earlier if a growth spurt or signs of puberty are evident and/or when there are plans for engaging in intense competitive sports or there is a family history of SCD (Maron, Seidman, & Seidman, 2004; Schwartz et al., 1996). (Level of Evidence: C)
4. Repeat TTE is recommended for the evaluation of patients with HCM with a change in clinical status or new cardiovascular event (Harris et al., 2006; Maron MS et al., "Effect," 2003; Maron et al., "Epidemiology," 2000; Dimitrow & Dubiel, 2005; Efthimiadis et al., 2009; Ommen, Shah, & Tajik, 2008; Sorajja et al., 2009). (Level of Evidence: B)

5. A transesophageal echocardiogram (TEE) is recommended for the intraoperative guidance of surgical myectomy (Grigg et al., 1992; Marwick et al., 1992; Yu et al., 2000). (Level of Evidence: B)
6. TTE or TEE with intracoronary contrast injection of the candidate's septal perforator(s) is recommended for the intraprocedural guidance of alcohol septal ablation (Sorajja et al., 2008; Faber et al., 2004; Monakier et al., 2004; Nagueh et al., 1998). (Level of Evidence: B)
7. TTE should be used to evaluate the effects of surgical myectomy or alcohol septal ablation for obstructive HCM (Ommen et al., 2005; Sorajja et al., 2008; Carasso et al., 2008; Fernandes et al., 2008; Jassal et al., 2006; Woo et al., 2005; Yoerger et al., 2006). (Level of Evidence: C)

Class IIa

1. TTE studies performed every 1 to 2 years can be useful in the serial evaluation of symptomatically stable patients with HCM to assess the degree of myocardial hypertrophy, dynamic obstruction, and myocardial function (Klues, Schiffers, & Maron, 1995; Wigle et al., 1995; Afonso et al., 2008). (Level of Evidence: C)
2. Exercise TTE can be useful in the detection and quantification of dynamic left ventricular outflow tract (LVOT) obstruction in the absence of resting outflow tract obstruction in patients with HCM (Maron et al., 2006; Maron MS et al., "Effect," 2003; Efthimiadis et al., 2009; Sorajja et al., 2009; Sherrid et al., 2005). (Level of Evidence: B)
3. TEE can be useful if TTE is inconclusive for clinical decision making about medical therapy and in situations such as planning for myectomy, exclusion of subaortic membrane or mitral regurgitation secondary to structural abnormalities of the mitral valve apparatus, or in assessment for the feasibility of alcohol septal ablation (Grigg et al., 1992; Marwick et al., 1992; Yu et al., 2000). (Level of Evidence: C)
4. TTE combined with the injection of an intravenous contrast agent is reasonable if the diagnosis of apical HCM or apical infarction or severity of hypertrophy is in doubt, particularly when other imaging modalities such as cardiovascular magnetic resonance (CMR) are not readily available, not diagnostic, or are contraindicated. (Level of Evidence: C)
5. Serial TTE studies are reasonable for clinically unaffected patients who have a first-degree relative with HCM when genetic status is unknown. Such follow-up may be considered every 12 to 18 months for children or adolescents from high-risk families and every 5 years for adult family members (Bos, Towbin, & Ackerman, 2009; Maron, Seidman, & Seidman, 2004; Hershberger et al., 2009; Schwartz et al., 1996). (Level of Evidence: C)

Class III: NO BENEFIT

1. TTE studies should not be performed more frequently than every 12 months in patients with HCM when it is unlikely that any changes have occurred that would have an impact on clinical decision making. (Level of Evidence: C)
2. Routine TEE and/or contrast echocardiography is not recommended when TTE images are diagnostic of HCM and/or there is no suspicion of fixed obstruction or intrinsic mitral valve pathology. (Level of Evidence: C)

Stress Testing

Class IIa

1. Treadmill exercise testing is reasonable to determine functional capacity and response to therapy in patients with HCM. (Level of Evidence: C)
2. Treadmill testing with monitoring of an ECG and blood pressure is reasonable for SCD risk stratification in patients with HCM (Sadoul et al., 1997; Olivetto et al., 1999; Ciampi et al., 2002). (Level of Evidence: B)
3. In patients with HCM who do not have a resting peak instantaneous gradient of greater than or equal to 50 mm Hg, exercise echocardiography is reasonable for the detection and quantification of exercise-induced dynamic LVOT obstruction (Maron et al., 2006; Frenneaux et al., 1990; Sadoul et al., 1997; Olivetto et al., 1999). (Level of Evidence: B)

Cardiac Magnetic Resonance

Class I

1. CMR imaging is indicated in patients with suspected HCM when echocardiography is inconclusive for diagnosis (Moon et al., 2004; Rickers et al., 2005). (Level of Evidence: B)
2. CMR imaging is indicated in patients with known HCM when additional information that may have an impact on management or decision making regarding invasive management, such as magnitude and distribution of hypertrophy or anatomy of the mitral valve apparatus or papillary muscles, is not adequately defined with echocardiography (Maron MS et al., "Hypertrophic," 2009; Moon et al., 2004; Rickers et al., 2005; Maron MS et al., "Prevalence," 2008; Maron et al., 2010). (Level of Evidence: B)

Class IIa

1. CMR imaging is reasonable in patients with HCM to define apical hypertrophy and/or aneurysm if echocardiography is inconclusive (Moon et al., 2004; Maron MS et al., "Prevalence," 2008) (Level of Evidence: B)

Class IIb

1. In selected patients with known HCM, when SCD risk stratification is inconclusive after documentation of the conventional risk factors (see Section 6.3.1 in the original guideline document), CMR imaging with assessment of late gadolinium enhancement (LGE) may be considered in resolving clinical decision making (Adabag et al., 2008; Maron MS et al., "Clinical," 2008; Rubinshtein et al., 2010; O'Hanlon et al., 2010; Moon et al., "Toward,"

2003). (Level of Evidence: C)

2. CMR imaging may be considered in patients with left ventricular (LV) hypertrophy and the suspicion of alternative diagnoses to HCM, including cardiac amyloidosis, Fabry disease, and genetic phenocopies such as LAMP2 cardiomyopathy (Gange, Link, & Maron, 2009; Maceira et al., 2005; Moon et al., "Gadolinium," 2003). (Level of Evidence: C)

Detection of Concomitant Coronary Disease

Class I

1. Coronary arteriography (invasive or computed tomographic imaging) is indicated in patients with HCM with chest discomfort who have an intermediate to high likelihood of coronary artery disease (CAD) when the identification of concomitant CAD will change management strategies. (Level of Evidence: C)

Class IIa

1. Assessment of coronary anatomy with computed tomographic angiography (CTA) is reasonable for patients with HCM with chest discomfort and a low likelihood of CAD to assess for possible concomitant CAD. (Level of Evidence: C)
2. Assessment of ischemia or perfusion abnormalities suggestive of CAD with single photon emission computed tomography (SPECT) or positron emission tomography (PET) myocardial perfusion imaging (MPI; because of excellent negative predictive value) is reasonable in patients with HCM with chest discomfort and a low likelihood of CAD to rule out possible concomitant CAD. (Level of Evidence: C)

Class III: NO BENEFIT

1. Routine SPECT MPI or stress echocardiography is not indicated for detection of "silent" CAD-related ischemia in patients with HCM who are asymptomatic. (Level of Evidence: C)
2. Assessment for the presence of blunted flow reserve (microvascular ischemia) using quantitative myocardial blood flow measurements by PET is not indicated for the assessment of prognosis in patients with HCM. (Level of Evidence: C)

Management of HCM

Asymptomatic Patients

Class I

1. For patients with HCM, it is recommended that comorbidities that may contribute to cardiovascular disease (e.g., hypertension, diabetes, hyperlipidemia, obesity) be treated in compliance with relevant existing guidelines (Redberg et al., 2009). (Level of Evidence: C)

Class IIa

1. Low-intensity aerobic exercise is reasonable as part of a healthy lifestyle for patients with HCM (Maron BJ et al., "American," 2003; Maron BJ et al., 2004). (Level of Evidence: C)

Class IIb

1. The usefulness of beta blockade and calcium channel blockers to alter clinical outcome is not well established for the management of asymptomatic patients with HCM with or without obstruction (Maron BJ et al., "American," 2003). (Level of Evidence: C)

Class III: HARM

1. Septal reduction therapy should not be performed for asymptomatic adult and pediatric patients with HCM with normal effort tolerance regardless of the severity of obstruction (Maron, 2002; Maron BJ et al., "American," 2003). (Level of Evidence: C)
2. In patients with HCM with resting or provokable outflow tract obstruction, regardless of symptom status, pure vasodilators and high-dose diuretics are potentially harmful (Braunwald et al., 1964; Maron, 2002). (Level of Evidence: C)

Symptomatic Patients

Pharmacologic Management

Class I

1. Beta-blocking drugs are recommended for the treatment of symptoms (angina or dyspnea) in adult patients with obstructive or nonobstructive HCM but should be used with caution in patients with sinus bradycardia or severe conduction disease (Braunwald et al., 1964; Maron, 2002; Maron BJ et al., "American," 2003; Spirito et al., 1997; Fifer & Vlahakes, 2008; Adelman et al., 1970; Cohen & Braunwald, 1967; Flamm, Harrison, & Hancock, 1968; Frank et al., 1978; Harrison et al., 1964; Stenson et al., 1973; Swanton et al., 1977; Wigle et al., 1974). (Level of Evidence: B)
2. If low doses of beta-blocking drugs are ineffective for controlling symptoms (angina or dyspnea) in patients with HCM, it is useful to titrate the dose to a resting heart rate of less than 60 to 65 bpm (up to generally accepted and recommended maximum doses of these drugs) (Braunwald et al., 1964;

Maron BJ et al., "American," 2003; Fifer & Vlahakes, 2008; Adelman et al., 1970; Cohen & Braunwald, 1967; Flamm, Harrison, & Hancock, 1968; Frank et al., 1978; Harrison et al., 1964; Stenson et al., 1973; Swanton, et al., 1977; Wigle et al., 1974). (Level of Evidence: B)

3. Verapamil therapy (starting in low doses and titrating up to 480 mg/d) is recommended for the treatment of symptoms (angina or dyspnea) in patients with obstructive or nonobstructive HCM who do not respond to beta-blocking drugs or who have side effects or contraindications to beta-blocking drugs. However, verapamil should be used with caution in patients with high gradients, advanced heart failure, or sinus bradycardia (Maron BJ et al., "American," 2003; Spirito et al., 1997; Fifer & Vlahakes, 2008; Bonow et al., 1981; Epstein & Rosing, 1981; Rosing et al., "Verapamil therapy: a new approach to the pharmacologic treatment of hypertrophic cardiomyopathy, II," 1979; Rosing et al., "Verapamil therapy: a new approach to the pharmacologic treatment of hypertrophic cardiomyopathy, I," 1979; Rosing et al., 1981). (Level of Evidence: B)
4. Intravenous phenylephrine (or another pure vasoconstricting agent) is recommended for the treatment of acute hypotension in patients with obstructive HCM who do not respond to fluid administration (Fifer & Vlahakes, 2008; Braunwald & Ebert, 1962; Wigle et al., 1965; Haley et al., 1999). (Level of Evidence: B)

Class IIa

1. It is reasonable to combine disopyramide with a beta-blocking drug or verapamil in the treatment of symptoms (angina or dyspnea) in patients with obstructive HCM who do not respond to beta-blocking drugs or verapamil alone (Maron BJ et al., "American," 2003; Spirito et al., 1997; Fifer & Vlahakes, 2008; Kimball, Bui, & Wigle, 1993; Pollick et al., 1988; Pollick, 1988; Sherrid, Delia, & Dwyer, 1988). (Level of Evidence: B)
2. It is reasonable to add oral diuretics in patients with nonobstructive HCM when dyspnea persists despite the use of beta blockers or verapamil or their combination (Wigle et al., 1995; Spirito et al., 1997). (Level of Evidence: C)

Class IIb

1. Beta-blocking drugs might be useful in the treatment of symptoms (angina or dyspnea) in children or adolescents with HCM, but patients treated with these drugs should be monitored for side effects, including depression, fatigue, or impaired scholastic performance. (Level of Evidence: C)
2. It may be reasonable to add oral diuretics with caution to patients with obstructive HCM when congestive symptoms persist despite the use of beta blockers or verapamil or their combination (Maron BJ et al., "American," 2003; Spirito et al., 1997; Fifer & Vlahakes, 2008). (Level of Evidence: C)
3. The usefulness of angiotensin-converting enzyme inhibitors or angiotensin receptor blockers in the treatment of symptoms (angina or dyspnea) in patients with HCM with preserved systolic function is not well established, and these drugs should be used cautiously (if at all) in patients with resting or provokable LVOT obstruction. (Level of Evidence: C)
4. In patients with HCM who do not tolerate verapamil or in whom verapamil is contraindicated, diltiazem may be considered. (Level of Evidence: C)

Class III: HARM

1. Nifedipine or other dihydropyridine calcium channel-blocking drugs are potentially harmful for treatment of symptoms (angina or dyspnea) in patients with HCM who have resting or provokable LVOT obstruction. (Level of Evidence: C)
2. Verapamil is potentially harmful in patients with obstructive HCM in the setting of systemic hypotension or severe dyspnea at rest. (Level of Evidence: C)
3. Digitalis is potentially harmful in the treatment of dyspnea in patients with HCM and in the absence of AF (Braunwald et al., 1964; Maron BJ et al., "American," 2003; Fifer & Vlahakes, 2008; Adelman et al., 1970; Braunwald et al., 1961; Braunwald, Brockenbrough, & Frye, 1962; Sonnenblick et al., 1966). (Level of Evidence: B)
4. The use of disopyramide alone without beta blockers or verapamil is potentially harmful in the treatment of symptoms (angina or dyspnea) in patients with HCM with AF because disopyramide may enhance atrioventricular conduction and increase the ventricular rate during episodes of AF (Maron BJ et al., "American," 2003; Wigle et al., 1985; Spirito et al., 1997; Bergfeldt, Schenck-Gustafsson, & Dahlqvist, 1992; Birkhead & Vaughan Williams, 1977; Jensen & Uhrenholt, 1976; Lara, Oakley, & Rowbotham, 1980; Morady, Scheinman, & Desai, 1982; Robertson & Miller, 1980). (Level of Evidence: B)
5. Dopamine, dobutamine, norepinephrine, and other intravenous positive inotropic drugs are potentially harmful for the treatment of acute hypotension in patients with obstructive HCM (Braunwald et al., 1964; Elesber et al., 2008; Braunwald & Ebert, 1962; Wigle et al., 1965; Haley et al., 1999; Krasnow et al., 1963; Pierce, Morrow, & Braunwald, 1964; Whalen et al., 1963). (Level of Evidence: B)

Invasive Therapies

Class I

1. Septal reduction therapy should be performed only by experienced operators* in the context of a comprehensive HCM clinical program and only for the treatment of eligible patients with severe drug-refractory symptoms and LVOT obstruction.† (van der Lee, et al., 2008) (Level of Evidence: C)

*Experienced operators are defined as an individual operator with a cumulative case volume of at least 20 procedures or an individual operator who is working in a dedicated HCM program with a cumulative total of at least 50 procedures (see Section 6.2.2.3 in the original guideline document).

†Eligible patients are defined by all of the following:

- a. Clinical: Severe dyspnea or chest pain (usually New York Heart Association [NYHA] functional classes III or IV) or occasionally other exertional symptoms (such as syncope or near syncope) that interfere with everyday activity or quality of life despite optimal medical therapy.
- b. Hemodynamic: Dynamic LVOT gradient at rest or with physiologic provocation ≥ 50 mm Hg associated with septal hypertrophy and systolic anterior

motion (SAM) of the mitral valve.

- c. Anatomic: Targeted anterior septal thickness sufficient to perform the procedure safely and effectively in the judgment of the individual operator.

Class IIa

1. Consultation with centers experienced in performing both surgical septal myectomy and alcohol septal ablation is reasonable when discussing treatment options for eligible patients with HCM with severe drug-refractory symptoms and LVOT obstruction. (Level of Evidence: C)
2. Surgical septal myectomy, when performed in experienced centers, can be beneficial and is the first consideration for the majority of eligible patients with HCM with severe drug-refractory symptoms and LVOT obstruction (Ommen et al., 2005; Sorajja et al., 2008; Woo et al., 2005; Firoozi et al., 2002; Ralph-Edwards et al., 2005; Smedira et al., 2008). (Level of Evidence: B)
3. Surgical septal myectomy, when performed at experienced centers, can be beneficial in symptomatic children with HCM and severe resting obstruction (>50 mm Hg) for whom standard medical therapy has failed (Theodoro et al., 1996). (Level of Evidence: C)
4. When surgery is contraindicated or the risk is considered unacceptable because of serious comorbidities or advanced age, alcohol septal ablation, when performed in experienced centers, can be beneficial in eligible adult patients with HCM with LVOT obstruction and severe drug-refractory symptoms (usually NYHA functional classes III or IV) (Sorajja et al., 2008; Fernandes et al., 2008; Gietzen et al., 2004; Kuhn et al., 2008; Kwon et al., 2008; Nagueh et al., 2001; Qin et al., 2001). (Level of Evidence: B)

Class IIb

1. Alcohol septal ablation, when performed in experienced centers, may be considered as an alternative to surgical myectomy for eligible adult patients with HCM with severe drug-refractory symptoms and LVOT obstruction when, after a balanced and thorough discussion, the patient expresses a preference for septal ablation (Fernandes et al., 2008; Firoozi et al., 2002; Kuhn et al., 2008; Nagueh et al., 2001; Qin et al., 2001). (Level of Evidence: B)
2. The effectiveness of alcohol septal ablation is uncertain in patients with HCM with marked (i.e., >30 mm) septal hypertrophy, and therefore the procedure is generally discouraged in such patients. (Level of Evidence: C)

Class III: HARM

1. Septal reduction therapy should not be done for adult patients with HCM who are asymptomatic with normal exercise tolerance or whose symptoms are controlled or minimized on optimal medical therapy. (Level of Evidence: C)
2. Septal reduction therapy should not be done unless performed as part of a program dedicated to the longitudinal and multidisciplinary care of patients with HCM. (Level of Evidence: C)
3. Mitral valve replacement for relief of LVOT obstruction should not be performed in patients with HCM in whom septal reduction therapy is an option. (Level of Evidence: C)
4. Alcohol septal ablation should not be done in patients with HCM with concomitant disease that independently warrants surgical correction (e.g., coronary artery bypass grafting for CAD, mitral valve repair for ruptured chordae) in whom surgical myectomy can be performed as part of the operation. (Level of Evidence: C)
5. Alcohol septal ablation should not be done in patients with HCM who are less than 21 years of age and is discouraged in adults less than 40 years of age if myectomy is a viable option. (Level of Evidence: C)

Pacing

Class IIa

1. In patients with HCM who have had a dual-chamber device implanted for non-HCM indications, it is reasonable to consider a trial of dual-chamber atrial-ventricular pacing (from the right ventricular apex) for the relief of symptoms attributable to LVOT obstruction (Erwin et al., 2000; Ommen et al., 1999; Slade et al., 1996; Gadler et al., 1999). (Level of Evidence: B)

Class IIb

1. Permanent pacing may be considered in medically refractory symptomatic patients with obstructive HCM who are suboptimal candidates for septal reduction therapy (Maron et al., 1999; Erwin et al., 2000; Ommen et al., 1999; Slade et al., 1996; Gadler et al., 1999). (Level of Evidence: B)

Class III: NO BENEFIT

1. Permanent pacemaker implantation for the purpose of reducing gradient should not be performed in patients with HCM who are asymptomatic or whose symptoms are medically controlled (Maron et al., 1999; Nishimura et al., 1997; Kappenberger et al., 1997). (Level of Evidence: C)
2. Permanent pacemaker implantation should not be performed as a first-line therapy to relieve symptoms in medically refractory symptomatic patients with HCM and LVOT obstruction who are candidates for septal reduction (Maron et al., 1999; Nishimura et al., 1997; Kappenberger et al., 1997). (Level of Evidence: B)

Patients with LV Systolic Dysfunction

Class I

1. Patients with nonobstructive HCM who develop systolic dysfunction with an ejection fraction (EF) less than or equal to 50% should be treated according to evidence-based medical therapy for adults with other forms of heart failure with reduced EF, including angiotensin-converting enzyme inhibitors, angiotensin receptor blockers, beta blockers, and other indicated drugs (Harris et al., 2006; Maron & Spirito, 1998). (Level of Evidence: B)
2. Other concomitant causes of systolic dysfunction (such as CAD) should be considered as potential contributors to systolic dysfunction in patients with HCM. (Level of Evidence: C)

Class IIb

1. ICD therapy may be considered in adult patients with advanced (as defined by NYHA functional class III or IV heart failure) nonobstructive HCM, on maximal medical therapy, and EF less than or equal to 50%, who do not otherwise have an indication for an ICD (Harris et al., 2006). (Level of Evidence: C)
2. For patients with HCM who develop systolic dysfunction, it may be reasonable to reassess the use of negative inotropic agents previously indicated, for example, verapamil, diltiazem, or disopyramide, and to consider discontinuing those therapies. (Level of Evidence: C)

Selection of Patients for Heart Transplantation

Class I

1. Patients with advanced heart failure (end stage*) and non-obstructive HCM not otherwise amenable to other treatment interventions, with EF less than or equal to 50% (or occasionally with preserved EF), should be considered for heart transplantation (Harris et al., 2006; Biagini et al., 2008). (Level of Evidence: B)
2. Symptomatic children with HCM with restrictive physiology who are not responsive to or appropriate candidates for other therapeutic interventions should be considered for heart transplantation (Gajarski et al., 2009; Towbin, 2002). (Level of Evidence: C)

*Characterized by systolic dysfunction (EF \leq 50%), often associated with left ventricular (LV) remodeling, including cavity enlargement and wall thinning, and because of diffuse myocardial scarring.

Class III: HARM

1. Heart transplantation should not be performed in mildly symptomatic patients of any age with HCM. (Level of Evidence: C)

Prevention of SCD

SCD Risk Stratification

Class I

1. All patients with HCM should undergo comprehensive SCD risk stratification at initial evaluation to determine the presence of the following: (Elliott et al., 2000; Maron BJ, "Contemporary," 2010; Elliott et al., 2006; Maron et al., 1981; Maron, "Risk," 2010; Cecchi, Maron, & Epstein, 1989; Elliott et al., 1999; Fananapazir et al., 1992; Maki et al., 1998; McKenna et al., 1981; Spirito et al., 2009) (Level of Evidence: B):
 - a. A personal history for ventricular fibrillation, sustained VT, or SCD events, including appropriate ICD therapy for ventricular tachyarrhythmias†
 - b. A family history for SCD events, including appropriate ICD therapy for ventricular tachyarrhythmias†
 - c. Unexplained syncope
 - d. Documented non-sustained ventricular tachycardia (NSVT) defined as 3 or more beats at greater than or equal to 120 bpm on ambulatory (Holter) ECG
 - e. Maximal LV wall thickness greater than or equal to 30 mm

†Appropriate ICD discharge is defined as ICD therapy triggered by VT or ventricular fibrillation, documented by stored intracardiac electrogram or cycle length data, in conjunction with the patient's symptoms immediately before and after device discharge.

Class IIa

1. It is reasonable to assess blood pressure response during exercise as part of SCD risk stratification in patients with HCM (Sadoul et al., 1997; Elliott et al., 2006; Maki et al., 1998). (Level of Evidence: B)
2. SCD risk stratification is reasonable on a periodic basis (every 12 to 24 months) for patients with HCM who have not undergone ICD implantation but would otherwise be eligible in the event that risk factors are identified (12 to 24 months). (Level of Evidence: C)

Class IIb

1. The usefulness of the following potential SCD risk modifiers is unclear but might be considered in selected patients with HCM for whom risk remains borderline after documentation of conventional risk factors:
 - a. CMR imaging with LGE (Adabag et al., 2006; Moon et al., "Toward," 2003) (Level of Evidence: C)
 - b. Double and compound mutations (i.e., >1) (Level of Evidence: C)
 - c. Marked LVOT obstruction (Maron MS et al., "Effect," 2003; Elliott et al., 2006; Efthimiadis et al., 2009; Maki et al., 1998) (Level of Evidence: B)

Class III: HARM

1. Invasive electrophysiologic testing as routine SCD risk stratification for patients with HCM should not be performed. (Level of Evidence: C)

Selection of Patients for ICDs

Class I

1. The decision to place an ICD in patients with HCM should include application of individual clinical judgment, as well as a thorough discussion of the strength of evidence, benefits, and risks to allow the informed patient's active participation in decision making (see Figure 4 in the original guideline document) (Maron BJ, "Contemporary," 2010; Maron BJ et al, "Efficacy," 2000; Maron BJ et al, "Implantable," 2007; Maron & Spirito, 2008). (Level of Evidence: C)
2. ICD placement is recommended for patients with HCM with prior documented cardiac arrest, ventricular fibrillation, or hemodynamically significant VT (Maron BJ et al, "Implantable," 2007; Cecchi, Maron, & Epstein, 1989; Elliott et al, 1999; Fananapazir et al, 1992). (Level of Evidence: B)

Class IIa

1. It is reasonable to recommend an ICD for patients with HCM with:
 - a. Sudden death presumably caused by HCM in 1 or more first-degree relatives (Bos et al, 2010). (Level of Evidence: C)
 - b. A maximum LV wall thickness greater than or equal to 30 mm (Elliott et al, 2000; Elliott et al, 2001; Spirito et al, 2000; Sorajja et al, 2006). (Level of Evidence: C)
 - c. One or more recent, unexplained syncopal episodes (Spirito et al, 2009). (Level of Evidence: C)
2. An ICD can be useful in select patients with NSVT (particularly those <30 years of age) in the presence of other SCD risk factors or modifiers‡ (Maron BJ, "Contemporary," 2010; Monserrat et al, 2003). (Level of Evidence: C)
3. An ICD can be useful in select patients with HCM with an abnormal blood pressure response with exercise in the presence of other SCD risk factors or modifiers‡ (Sadoul et al, 1997; Olivotto et al, 1999; Maki et al, 1998). (Level of Evidence: C)
4. It is reasonable to recommend an ICD for high-risk children with HCM, based on unexplained syncope, massive LV hypertrophy, or family history of SCD, after taking into account the relatively high complication rate of long-term ICD implantation. (Level of Evidence: C)

‡SCD risk modifiers are discussed in Section 6.3.1.2 in the original guideline document.

Class IIb

1. The usefulness of an ICD is uncertain in patients with HCM with isolated bursts of NSVT when in the absence of any other SCD risk factors or modifiers‡ (Maron, 2010). (Level of Evidence: C)
2. The usefulness of an ICD is uncertain in patients with HCM with an abnormal blood pressure response with exercise when in the absence of any other SCD risk factors or modifiers‡, particularly in the presence of significant outflow obstruction (Sadoul et al, 1997; Olivotto et al, 1999; Maki et al, 1998). (Level of Evidence: C)

‡SCD risk modifiers are discussed in Section 6.3.1.2 in the original guideline document.

Class III: HARM

1. ICD placement as a routine strategy in patients with HCM without an indication of increased risk is potentially harmful. (Level of Evidence: C)
2. ICD placement as a strategy to permit patients with HCM to participate in competitive athletics is potentially harmful. (Level of Evidence: C)
3. ICD placement in patients who have an identified HCM genotype in the absence of clinical manifestations of HCM is potentially harmful. (Level of Evidence: C)

Selection of ICD Device Type

Class IIa

1. In patients with HCM who meet indications for ICD implantation, single-chamber devices are reasonable in younger patients without a need for atrial or ventricular pacing (Hauser et al, 2008; Boriani et al, 2004; Kleemann et al, 2007; Maisel et al, 2001). (Level of Evidence: B)
2. In patients with HCM who meet indications for ICD implantation, dual-chamber ICDs are reasonable for patients with sinus bradycardia and/or paroxysmal AF (Boriani et al, 2004). (Level of Evidence: C)
3. In patients with HCM who meet indications for ICD implantation, dual-chamber ICDs are reasonable for patients with elevated resting outflow gradients greater than 50 mm Hg and significant heart failure symptoms who may benefit from right ventricular pacing (most commonly, but not limited to, patients >65 years of age) (Maron et al, 1999; Nishimura et al, 1997; Kappenberger et al, 1997; Boriani et al, 2004). (Level of Evidence: B)

Participation in Competitive or Recreational Sports and Physical Activity

Class IIa

1. It is reasonable for patients with HCM to participate in low-intensity competitive sports (e.g., golf and bowling) (Maron et al, 2005; Pelliccia et al,

2005). (Level of Evidence: C)

2. It is reasonable for patients with HCM to participate in a range of recreational sporting activities as outlined in Table 4 in the original guideline document (Maron et al., 2004). (Level of Evidence: C)

Class III: HARM

1. Patients with HCM should not participate in intense competitive sports regardless of age, sex, race, presence or absence of LVOT obstruction, prior septal reduction therapy, or implantation of a cardioverter-defibrillator for high-risk status (Maron, 2003; Maron BJ et al., "Sudden," 2009; Maron BJ et al., 2005; Pelliccia et al., 2005; Maron, Epstein, & Roberts, 1986; Maron BJ et al., "Relationship," 2003; Corrado et al., 2003). (Level of Evidence: C)

Management of AF

Class I

1. Anticoagulation with vitamin K antagonists (i.e., warfarin, to an international normalized ratio of 2.0 to 3.0) is indicated in patients with paroxysmal, persistent, or chronic AF and HCM (Olivetto et al., 2001; Fuster et al., 2011; Maron et al., 2002). (Anticoagulation with direct thrombin inhibitors [i.e., dabigatran] may represent another option to reduce the risk of thromboembolic events, but data for patients with HCM are not available) (Connolly et al., 2009). (Level of Evidence: C)
2. Ventricular rate control in patients with HCM with AF is indicated for rapid ventricular rates and can require high doses of beta antagonists and non-dihydropyridine calcium channel blockers (Olivetto et al., 2001; Fuster et al., 2011). (Level of Evidence: C)

§Dabigatran should not be used in patients with prosthetic valves, hemodynamically significant valve disease, advanced liver failure, or severe renal failure (creatinine clearance <15 mL/min [Connolly et al., 2009]).

Class IIa

1. Disopyramide (with ventricular rate-controlling agents) and amiodarone are reasonable antiarrhythmic agents for AF in patients with HCM (Fuster et al., 2011; Tendra et al., 1993). (Level of Evidence: B)
2. Radiofrequency ablation for AF can be beneficial in patients with HCM who have refractory symptoms or who are unable to take antiarrhythmic drugs (Bunch et al., 2008; Gaita et al., 2007; Kilicaslan et al., 2006; Callans 2008; Di Donna et al., 2010). (Level of Evidence: B)
3. Maze procedure with closure of left atrial (LA) appendage is reasonable in patients with HCM with a history of AF, either during septal myectomy or as an isolated procedure in selected patients. (Level of Evidence: C)

Class IIb

1. Sotalol, dofetilide, and dronedarone might be considered alternative antiarrhythmic agents in patients with HCM, especially in those with an ICD, but clinical experience is limited. (Level of Evidence: C)

Other Issues

Pregnancy/Delivery

Class I

1. In women with HCM who are asymptomatic or whose symptoms are controlled with beta-blocking drugs, the drugs should be continued during pregnancy, but increased surveillance for fetal bradycardia or other complications is warranted (Bos, Towbin, & Ackerman, 2009; Hersherberger et al., 2009; Bascou et al., 1993; Fitzgerald-Butt et al., 2010). (Level of Evidence: C)
2. For patients (mother or father) with HCM, genetic counseling is indicated before planned conception. (Level of Evidence: C)
3. In women with HCM and resting or provokable LVOT obstruction greater than or equal to 50 mm Hg and/or cardiac symptoms not controlled by medical therapy alone, pregnancy is associated with increased risk, and these patients should be referred to a high-risk obstetrician. (Level of Evidence: C)
4. The diagnosis of HCM among asymptomatic women is not considered a contraindication for pregnancy, but patients should be carefully evaluated in regard to the risk of pregnancy. (Level of Evidence: C)

Class IIa

1. For women with HCM whose symptoms are controlled (mild to moderate), pregnancy is reasonable, but expert maternal/fetal medical specialist care, including cardiovascular and prenatal monitoring, is advised. (Level of Evidence: C)

Class III: HARM

1. For women with advanced heart failure symptoms and HCM, pregnancy is associated with excess morbidity/mortality. (Level of Evidence: C)

Definitions:

		Size of Treatment Effect					
		CLASS I	CLASS IIa	CLASS IIb	CLASS III <i>No Benefit</i> or Class III <i>Harm</i>		
		<i>Benefit >>> Risk</i>	<i>Benefit >> Risk</i> <i>Additional studies with focused objectives needed</i>	<i>Benefit ≥ Risk</i> <i>Additional studies with broad objectives needed; additional registry data would be helpful</i>		Procedure/Test	Treatment
		Procedure/Treatment SHOULD be performed/ administered	IT IS REASONABLE to perform procedure/administer treatment	Procedure/Treatment MAY BE CONSIDERED	COR III: No Benefit	Not helpful	No proven benefit
					COR III: Harm	Excess Cost without Benefit or Harmful	Harmful to Patients
Estimate of Certainty (Precision) of Treatment Effect	LEVEL A	<ul style="list-style-type: none">Recommendation that procedure or treatment is useful/effectiveSufficient evidence from multiple randomized trials or meta-analyses	<ul style="list-style-type: none">Recommendation in favor of treatment or procedure being useful/effectiveSome conflicting evidence from multiple randomized trials or meta-analyses	<ul style="list-style-type: none">Recommendation's usefulness/efficacy less well establishedGreater conflicting evidence from multiple randomized trials or meta-analyses	<ul style="list-style-type: none">Recommendation that procedure or treatment is not useful/effective and may be harmfulSufficient evidence from multiple randomized trials or meta-analyses		
	LEVEL B	<ul style="list-style-type: none">Recommendation that procedure or treatment is useful/effectiveEvidence from single randomized trial or nonrandomized studies	<ul style="list-style-type: none">Recommendation in favor of treatment or procedure being useful/effectiveSome conflicting evidence from single randomized trial or nonrandomized studies	<ul style="list-style-type: none">Recommendation's usefulness/efficacy less well establishedGreater conflicting evidence from single randomized trial or nonrandomized studies	<ul style="list-style-type: none">Recommendation that procedure or treatment is not useful/effective and may be harmfulEvidence from single randomized trial or nonrandomized studies		
	LEVEL C	<ul style="list-style-type: none">Recommendation that procedure or treatment is useful/effectiveOnly expert opinion, case studies, or standard of care	<ul style="list-style-type: none">Recommendation in favor of treatment or procedure being useful/effectiveOnly diverging expert opinion, case studies, or standard of care	<ul style="list-style-type: none">Recommendation's usefulness/efficacy less well establishedOnly diverging expert opinion, case studies, or standard of care	<ul style="list-style-type: none">Recommendation that procedure or treatment is not useful/effective and may be harmfulOnly expert opinion, case studies, or standard of care		

A recommendation with Level of Evidence B or C does not imply that the recommendation is weak. Many important clinical questions addressed in the

guidelines do not lend themselves to clinical trials. Although randomized trials are unavailable, there may be a very clear clinical consensus that a particular test or therapy is useful or effective.

*Data available from clinical trials or registries about the usefulness/efficacy in different subpopulations, such as gender, age, history of diabetes, history of prior myocardial infarction, history of heart failure, and prior aspirin use. A recommendation with Level of Evidence B or C does not imply that the recommendation is weak. Many important clinical questions addressed in the guidelines do not lend themselves to clinical trials. Even though randomized trials are not available, there may be a very clear clinical consensus that a particular test or therapy is useful or effective.

Clinical Algorithm(s)

The following algorithms are provided in the original guideline document:

- Treatment Algorithm
- Indications for ICDs in HCM
- Management of AF in HCM

Scope

Disease/Condition(s)

Hypertrophic cardiomyopathy

Guideline Category

Diagnosis

Evaluation

Management

Prevention

Risk Assessment

Treatment

Clinical Specialty

Cardiology

Family Practice

Internal Medicine

Preventive Medicine

Intended Users

Physicians

Guideline Objective(s)

To assist healthcare providers in clinical decision making by describing a range of generally acceptable approaches for the diagnosis, management, and prevention of specific diseases or conditions

Target Population

Interventions and Practices Considered

Diagnosis/Evaluation

1. Genetic testing and family screening for hypertrophic cardiomyopathy (HCM)
2. Electrocardiogram (ECG) (12 lead ECG, 24-hour ambulatory monitoring)
3. Echocardiography (transthoracic)
4. Stress testing
5. Cardiac magnetic resonance imaging
6. Coronary arteriography (invasive or computer tomographic)
7. Single photon emission computed tomography (SPECT) or positron emission tomography (PET) myocardial perfusion imaging
8. Sudden cardiac death (SCD) risk stratification

Management/Treatment

1. Management of comorbid conditions in asymptomatic patients
2. Pharmacological agents
 - Beta blockers
 - Verapamil (diltiazem if verapamil is contraindicated)
 - Intravenous phenylephrine
 - Disopyramide with a beta-blocking drug or verapamil
 - Oral diuretics
3. Invasive therapies
 - Septal reduction therapy
 - Surgical septal myectomy
 - Alcohol septal ablation
4. Pacing
5. Management of patients with left ventricular systolic dysfunction (angiotensin-converting enzyme inhibitors, angiotensin receptor blockers, beta blockers, other indicated drugs, implantable cardioverter-defibrillator [ICD] in certain patients)
6. Selection of patients for heart transplantation
7. Selection of patients for ICD placement
8. Selection of ICD device type
9. Patient participation in sports or physical activity
10. Management of atrial fibrillation (AF)
 - Anticoagulation (vitamin K antagonists, thrombin inhibitors)
 - Beta antagonists and non-dihydropyridine calcium channel blockers for ventricular rate control
 - Disopyramide (with ventricular rate-controlling agents) and amiodarone (sotalol, dofetilide, dronedarone may be alternatives)
 - Radiofrequency ablation for AF
 - Maze procedure with closure of left atrial appendage
11. Management of pregnancy and delivery in women with HCM

Major Outcomes Considered

- Mortality
- Morbidity
- Risk of cardiac events
- Symptom control
- Adverse events associated with treatment

Methodology

Methods Used to Collect/Select the Evidence

Hand-searches of Published Literature (Primary Sources)

Description of Methods Used to Collect/Select the Evidence

An extensive evidence review was conducted through January 2011. Searches were limited to studies, reviews, and other evidence conducted in human subjects and published in English. Key search words included, but were not limited to, hypertrophic cardiomyopathy (HCM), surgical myectomy, ablation, exercise, sudden cardiac death (SCD), athletes, dual-chamber pacing, left ventricular outflow tract (LVOT) obstruction, alcohol septal ablation, automobile driving and implantable cardioverter-defibrillators (ICDs), catheter ablation, defibrillators, genetics, genotype, medical management, magnetic resonance imaging, pacing, permanent pacing, phenotype, pregnancy, risk stratification, sudden death in athletes, surgical septal myectomy, and septal reduction. Additionally, the committee reviewed documents related to the subject matter previously published by the American College of Cardiology Foundation (ACCF) and the American Heart Association (AHA).

Number of Source Documents

Not stated

Methods Used to Assess the Quality and Strength of the Evidence

Weighting According to a Rating Scheme (Scheme Given)

Rating Scheme for the Strength of the Evidence

Applying Classification of Recommendations and Level of Evidence

		Size of Treatment Effect					
		CLASS I <i>Benefit >>> Risk</i> Procedure/Treatment SHOULD be performed/ administered	CLASS IIa <i>Benefit >> Risk</i> <i>Additional studies with focused objectives needed</i> IT IS REASONABLE to perform procedure/administer treatment	CLASS IIb <i>Benefit ≥ Risk</i> <i>Additional studies with broad objectives needed; additional registry data would be helpful</i> Procedure/Treatment MAY BE CONSIDERED	CLASS III <i>No Benefit</i> or Class III <i>Harm</i>		
						Procedure/Test	Treatment
					COR III: No Benefit	Not helpful	No proven benefit
					COR III: Harm	Excess Cost without Benefit or Harmful	Harmful to Patients
Estimate of Certainty (Precision) of Treatment Effect	LEVEL A Multiple populations evaluated* Data derived from multiple randomized clinical trials or meta-analyses	<ul style="list-style-type: none">Recommendation that procedure or treatment is useful/effectiveSufficient evidence from multiple randomized trials or meta-analyses	<ul style="list-style-type: none">Recommendation in favor of treatment or procedure being useful/effectiveSome conflicting evidence from multiple randomized trials or meta-analyses	<ul style="list-style-type: none">Recommendation's usefulness/efficacy less well establishedGreater conflicting evidence from multiple randomized trials or meta-analyses	<ul style="list-style-type: none">Recommendation that procedure or treatment is not useful/effective and may be harmfulSufficient evidence from multiple randomized trials or meta-analyses		
	LEVEL B Limited populations evaluated*	<ul style="list-style-type: none">Recommendation that procedure or treatment is useful/effectiveEvidence from	<ul style="list-style-type: none">Recommendation in favor of treatment or procedure being useful/effective	<ul style="list-style-type: none">Recommendation's usefulness/efficacy less well establishedGreater conflicting	<ul style="list-style-type: none">Recommendation that procedure or treatment is not useful/effective and may be harmfulEvidence from single randomized		

	Data derived from a single randomized clinical trials or nonrandomized studies	Size of Treatment Effect from single randomized trial or nonrandomized studies	• Some conflicting evidence from single randomized trial or nonrandomized studies	evidence from single randomized trial or nonrandomized studies	trial or nonrandomized studies
	<p>LEVEL C</p> <p>Very limited populations evaluated*</p> <p>Only consensus opinion of experts, case studies or standard of care</p>	<ul style="list-style-type: none"> • Recommendation that procedure or treatment is useful/effective • Only expert opinion, case studies, or standard of care 	<ul style="list-style-type: none"> • Recommendation in favor of treatment or procedure being useful/effective • Only diverging expert opinion, case studies, or standard of care 	<ul style="list-style-type: none"> • Recommendation's usefulness/efficacy less well established • Only diverging expert opinion, case studies, or standard of care 	<ul style="list-style-type: none"> • Recommendation that procedure or treatment is not useful/effective and may be harmful • Only expert opinion, case studies, or standard of care

A recommendation with Level of Evidence B or C does not imply that the recommendation is weak. Many important clinical questions addressed in the guidelines do not lend themselves to clinical trials. Although randomized trials are unavailable, there may be a very clear clinical consensus that a particular test or therapy is useful or effective.

*Data available from clinical trials or registries about the usefulness/efficacy in different subpopulations, such as gender, age, history of diabetes, history of prior myocardial infarction, history of heart failure, and prior aspirin use. A recommendation with Level of Evidence B or C does not imply that the recommendation is weak. Many important clinical questions addressed in the guidelines do not lend themselves to clinical trials. Even though randomized trials are not available, there may be a very clear clinical consensus that a particular test or therapy is useful or effective.

Methods Used to Analyze the Evidence

Review of Published Meta-Analyses

Systematic Review with Evidence Tables

Description of the Methods Used to Analyze the Evidence

To provide clinicians with a comprehensive set of data, whenever deemed appropriate or when published, the absolute risk difference and number needed to treat or harm are provided in the guideline, along with confidence intervals and data related to the relative treatment effects, such as odds ratio, relative risk, hazard ratio, or incidence rate ratio.

In analyzing the data and developing the recommendations and supporting text, the writing committee used evidence based methodologies developed by the Task Force. The committee reviewed and ranked evidence supporting current recommendations with the weight of evidence ranked as Level A if the data were derived from multiple randomized clinical trials (RCTs) or meta-analyses. The committee ranked available evidence as Level B when data were derived from a single RCT or nonrandomized studies. Evidence was ranked as Level C when the primary source of the recommendation was consensus opinion, case studies, or standard of care. In the narrative portions of these guidelines, evidence is generally presented in chronological order of development. Studies are identified as observational, retrospective, prospective, or randomized when appropriate. For certain conditions for which inadequate data are available, recommendations are based on expert consensus and clinical experience and ranked as Level C. An example is the use of penicillin for pneumococcal pneumonia, for which there are no RCTs and treatment is based on clinical experience. When recommendations at Level C are supported by historical clinical data, appropriate references (including clinical reviews) are cited if available. For issues where sparse data are available, a survey of current practice among the clinicians on the writing committee was the basis for Level C recommendations and no references are cited. The schema for Classification of Recommendations and Level of Evidence is summarized in Table 1 (see the "Rating Scheme for the Strength of the Evidence" field), which also illustrates how the grading system provides an estimate of the size and the certainty of the treatment effect. A new addition to the ACCF/AHA methodology is separation of the Class III recommendations to delineate whether the recommendation is determined to be of "no benefit" or associated with "harm" to the patient. In addition, in view of the increasing number of comparative effectiveness studies, comparator verbs and suggested phrases for writing recommendations for the comparative effectiveness of one treatment/strategy with respect to another for Class of Recommendation I and IIa, Level of Evidence A or B only have been added.

Methods Used to Formulate the Recommendations

Expert Consensus

Description of Methods Used to Formulate the Recommendations

Experts in the subject under consideration have been selected from both organizations to examine subject-specific data and write guidelines in partnership with representatives from other medical practitioner and specialty groups. Writing committees are specifically charged to perform a formal literature review, weigh the strength of evidence for or against particular tests, treatments, or procedures, and include estimates of expected health outcomes where data exist. Patient-specific modifiers, comorbidities, and issues of patient preference that may influence the choice of tests or therapies are considered. When available, information from studies on cost is considered, but data on efficacy and clinical outcomes constitute the primary basis for recommendations in these guidelines.

The recommendations listed in the original guideline document are, whenever possible, evidence based.

The committee was composed of physicians and cardiac surgeons with expertise in hypertrophic cardiomyopathy (HCM), invasive cardiology, non-invasive testing and imaging, pediatric cardiology, electrophysiology, and genetics. The committee included representatives from the American Association for Thoracic Surgery, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons.

Rating Scheme for the Strength of the Recommendations

See the "Rating Scheme for the Strength of the Evidence" field, above.

Cost Analysis

A formal cost analysis was not performed and published cost analyses were not reviewed.

Method of Guideline Validation

External Peer Review

Internal Peer Review

Description of Method of Guideline Validation

This document was reviewed by 2 outside reviewers nominated by both the American College of Cardiology Foundation (ACCF) and the American Heart Association (AHA), as well as 2 reviewers each from the American Association for Thoracic Surgery, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, and the Society of Thoracic Surgeons. Other content reviewers included members from the ACCF Adult Congenital and Pediatric Cardiology Council, ACCF Surgeons' Scientific Council, and ACCF Interventional Scientific Council. All information on reviewers' relationship with industry (RWI) was distributed to the writing committee and is published in this document (see Appendix 2 in the original guideline document).

This document was approved for publication by the governing bodies of the ACCF and the AHA and endorsed by the American Association for Thoracic Surgery, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons.

Evidence Supporting the Recommendations

References Supporting the Recommendations

Adabag AS, Kuskowski MA, Maron BJ. Determinants for clinical diagnosis of hypertrophic cardiomyopathy. *Am J Cardiol*. 2006 Dec 1;98(11):1507-11. [PubMed](#)

Adabag AS, Maron BJ, Appelbaum E, Harrigan CJ, Buross JL, Gibson CM, Lesser JR, Hanna CA, Udelson JE, Manning WJ, Maron MS. Occurrence

and frequency of arrhythmias in hypertrophic cardiomyopathy in relation to delayed enhancement on cardiovascular magnetic resonance. *J Am Coll Cardiol*. 2008 Apr 8;51(14):1369-74. [PubMed](#)

Adelman AG, Shah PM, Gramiak R, Wigle ED. Long-term propranolol therapy in muscular subaortic stenosis. *Br Heart J*. 1970 Nov;32(6):804-11. [PubMed](#)

Afonso LC, Bernal J, Bax JJ, Abraham TP. Echocardiography in hypertrophic cardiomyopathy: the role of conventional and emerging technologies. *JACC Cardiovasc Imaging*. 2008 Nov;1(6):787-800. [85 references] [PubMed](#)

Andersen PS, Havndrup O, Hougs L, Sorensen KM, Jensen M, Larsen LA, Hedley P, Thomsen AR, Moolman-Smook J, Christiansen M, Bundgaard H. Diagnostic yield, interpretation, and clinical utility of mutation screening of sarcomere encoding genes in Danish hypertrophic cardiomyopathy patients and relatives. *Hum Mutat*. 2009 Mar;30(3):363-70. [PubMed](#)

Arad M, Maron BJ, Gorham JM, Johnson WH Jr, Saul JP, Perez-Atayde AR, Spirito P, Wright GB, Kanter RJ, Seidman CE, Seidman JG. Glycogen storage diseases presenting as hypertrophic cardiomyopathy. *N Engl J Med*. 2005 Jan 27;352(4):362-72. [PubMed](#)

Bascou V, Ferrandis J, Bauer V, Bouret JM, de Meeus JB, Magnin G. [Obstructive myocardiopathy and pregnancy]. *J Gynecol Obstet Biol Reprod (Paris)*. 1993;22(3):309-11. [13 references] [PubMed](#)

Bergfeldt L, Schenck-Gustafsson K, Dahlqvist R. Comparative class 1 electrophysiologic and anticholinergic effects of disopyramide and its main metabolite (mono-N-dealkylated disopyramide) in healthy humans. *Cardiovasc Drugs Ther*. 1992 Oct;6(5):529-37. [PubMed](#)

Biagini E, Spirito P, Leone O, Picchio FM, Cocco F, Ragni L, Lofiego C, Grigioni F, Potena L, Rocchi G, Bacchi-Reggiani L, Boriani G, Prandstraller D, Arbustini E, Branzi A, Rapezzi C. Heart transplantation in hypertrophic cardiomyopathy. *Am J Cardiol*. 2008 Feb 1;101(3):387-92. [PubMed](#)

Binder J, Ommen SR, Gersh BJ, Van Driest SL, Tajik AJ, Nishimura RA, Ackerman MJ. Echocardiography-guided genetic testing in hypertrophic cardiomyopathy: septal morphological features predict the presence of myofilament mutations. *Mayo Clin Proc*. 2006 Apr;81(4):459-67. [PubMed](#)

Birkhead JS, Vaughan Williams EM. Dual effect of disopyramide on atrial and atrioventricular conduction and refractory periods. *Br Heart J*. 1977 Jun;39(6):657-60. [PubMed](#)

Bonow RO, Rosing DR, Bacharach SL, Green MV, Kent KM, Lipson LC, Maron BJ, Leon MB, Epstein SE. Effects of verapamil on left ventricular systolic function and diastolic filling in patients with hypertrophic cardiomyopathy. *Circulation*. 1981 Oct;64(4):787-96. [PubMed](#)

Boriani G, Maron BJ, Shen WK, Spirito P. Prevention of sudden death in hypertrophic cardiomyopathy: but which defibrillator for which patient. *Circulation*. 2004 Oct 12;110(15):e438-42. [17 references] [PubMed](#)

Bos JM, Maron BJ, Ackerman MJ, Haas TS, Sorajja P, Nishimura RA, Gersh BJ, Ommen SR. Role of family history of sudden death in risk stratification and prevention of sudden death with implantable defibrillators in hypertrophic cardiomyopathy. *Am J Cardiol*. 2010 Nov 15;106(10):1481-6. [PubMed](#)

Bos JM, Towbin JA, Ackerman MJ. Diagnostic, prognostic, and therapeutic implications of genetic testing for hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2009 Jul 14;54(3):201-11. [112 references] [PubMed](#)

Braunwald E, Bloodwell RD, Goldberg LI, Morrow AG. Studies on digitalis. IV. Observations in man on the effects of digitalis preparations on the contractility of the non-failing heart and on total vascular resistance. *J Clin Invest*. 1961 Jan;40(1):52-9. [PubMed](#)

Braunwald E, Brockenbrough EC, Frye RL. Studies on digitalis. V. Comparison of the effects of ouabain on left ventricular dynamics in valvular aortic stenosis and hypertrophic subaortic stenosis. *Circulation*. 1962 Aug;26:166-73. [PubMed](#)

Braunwald E, Ebert PA. Hemodynamic alterations in idiopathic hypertrophic subaortic stenosis induced by sympathomimetic drugs. *Am J Cardiol.* 1962 Oct;10:489-95. [PubMed](#)

Braunwald E, Lambrew CT, Rockoff SD, Ross J Jr, Morrow AG. Idiopathic hypertrophic subaortic stenosis. I. A description of the disease based upon an analysis of 64 patients. *Circulation.* 1964 Nov;30(Suppl 4):3-119. [PubMed](#)

Bunch TJ, Munger TM, Friedman PA, Asirvatham SJ, Brady PA, Cha YM, Rea RF, Shen WK, Powell BD, Ommen SR, Monahan KH, Haroldson JM, Packer DL. Substrate and procedural predictors of outcomes after catheter ablation for atrial fibrillation in patients with hypertrophic cardiomyopathy. *J Cardiovasc Electrophysiol.* 2008 Oct;19(10):1009-14. [PubMed](#)

Callans DJ. Ablation of atrial fibrillation in the setting of hypertrophic cardiomyopathy. *J Cardiovasc Electrophysiol.* 2008 Oct;19(10):1015-6. [PubMed](#)

Carasso S, Woo A, Yang H, Schwartz L, Vannan MA, Jamorski M, Linghorne M, Wigle ED, Rakowski H. Myocardial mechanics explains the time course of benefit for septal ethanol ablation for hypertrophic cardiomyopathy. *J Am Soc Echocardiogr.* 2008 May;21(5):493-9. [PubMed](#)

Cecchi F, Maron BJ, Epstein SE. Long-term outcome of patients with hypertrophic cardiomyopathy successfully resuscitated after cardiac arrest. *J Am Coll Cardiol.* 1989 May;13(6):1283-8. [PubMed](#)

Christiaans I, Birnie E, van Langen IM, van Spaendonck-Zwarts KY, van Tintelen JP, van den Berg MP, Atsma DE, Helderman-van den Enden AT, Pinto YM, Hermans-van Ast JF, Bonsel GJ, Wilde AA. The yield of risk stratification for sudden cardiac death in hypertrophic cardiomyopathy myosin-binding protein C gene mutation carriers: focus on predictive screening. *Eur Heart J.* 2010 Apr;31(7):842-8. [PubMed](#)

Christiaans I, Lekanne dit Deprez RH, van Langen IM, Wilde AA. Ventricular fibrillation in MYH7-related hypertrophic cardiomyopathy before onset of ventricular hypertrophy. *Heart Rhythm.* 2009 Sep;6(9):1366-9. [PubMed](#)

Christiaans I, van Langen IM, Birnie E, Bonsel GJ, Wilde AA, Smets EM. Genetic counseling and cardiac care in predictively tested hypertrophic cardiomyopathy mutation carriers: the patients' perspective. *Am J Med Genet A.* 2009 Jul;149A(7):1444-51. [PubMed](#)

Christiaans I, van Langen IM, Birnie E, Bonsel GJ, Wilde AA, Smets EM. Quality of life and psychological distress in hypertrophic cardiomyopathy mutation carriers: a cross-sectional cohort study. *Am J Med Genet A.* 2009 Feb 15;149A(4):602-12. [PubMed](#)

Ciampi Q, Betocchi S, Lombardi R, Manganelli F, Storto G, Losi MA, Pezzella E, Finizio F, Cuocolo A, Chiariello M. Hemodynamic determinants of exercise-induced abnormal blood pressure response in hypertrophic cardiomyopathy. *J Am Coll Cardiol.* 2002 Jul 17;40(2):278-84. [PubMed](#)

Cohen LS, Braunwald E. Amelioration of angina pectoris in idiopathic hypertrophic subaortic stenosis with beta-adrenergic blockade. *Circulation.* 1967 May;35(5):847-51. [PubMed](#)

Connolly SJ, Ezekowitz MD, Yusuf S, Eikelboom J, Oldgren J, Parekh A, Pogue J, Reilly PA, Themeles E, Varrone J, Wang S, Alings M, Xavier D, Zhu J, Diaz R, Lewis BS, Darius H, Diener HC, Joyner CD, Wallentin L, RE-LY Steering Committee and Investigators. Dabigatran versus warfarin in patients with atrial fibrillation. *N Engl J Med.* 2009 Sep 17;361(12):1139-51. [PubMed](#)

Corrado D, Basso C, Rizzoli G, Schiavon M, Thiene G. Does sports activity enhance the risk of sudden death in adolescents and young adults. *J Am Coll Cardiol.* 2003 Dec 3;42(11):1959-63. [PubMed](#)

Di Donna P, Olivetto I, Delcre SD, Caponi D, Scaglione M, Nault I, Montefusco A, Girolami F, Cecchi F, Haissaguerre M, Gaita F. Efficacy of catheter ablation for atrial fibrillation in hypertrophic cardiomyopathy: impact of age, atrial remodelling, and disease progression. *Europace.* 2010 Mar;12(3):347-55. [PubMed](#)

Dimitrow PP, Dubiel JS. Echocardiographic risk factors predisposing to sudden cardiac death in hypertrophic cardiomyopathy. *Heart*. 2005 Jan;91(1):93-4. [PubMed](#)

Effthimiadis GK, Parcharidou DG, Giannakoulas G, Pagourelas ED, Charalampidis P, Savopoulos G, Ziakas A, Karvounis H, Styliadis IH, Parcharidis GE. Left ventricular outflow tract obstruction as a risk factor for sudden cardiac death in hypertrophic cardiomyopathy. *Am J Cardiol*. 2009 Sep 1;104(5):695-9. [PubMed](#)

Elesber A, Nishimura RA, Rihal CS, Ommen SR, Schaff HV, Holmes DR Jr. Utility of isoproterenol to provoke outflow tract gradients in patients with hypertrophic cardiomyopathy. *Am J Cardiol*. 2008 Feb 15;101(4):516-20. [PubMed](#)

Elliott PM, Gimeno Blanes JR, Mahon NG, Poloniecki JD, McKenna WJ. Relation between severity of left-ventricular hypertrophy and prognosis in patients with hypertrophic cardiomyopathy. *Lancet*. 2001 Feb 10;357(9254):420-4. [PubMed](#)

Elliott PM, Gimeno JR, Tome MT, Shah J, Ward D, Thaman R, Mogensen J, McKenna WJ. Left ventricular outflow tract obstruction and sudden death risk in patients with hypertrophic cardiomyopathy. *Eur Heart J*. 2006 Aug;27(16):1933-41. [PubMed](#)

Elliott PM, Poloniecki J, Dickie S, Sharma S, Monserrat L, Varnava A, Mahon NG, McKenna WJ. Sudden death in hypertrophic cardiomyopathy: identification of high risk patients. *J Am Coll Cardiol*. 2000 Dec;36(7):2212-8. [PubMed](#)

Elliott PM, Sharma S, Varnava A, Poloniecki J, Rowland E, McKenna WJ. Survival after cardiac arrest or sustained ventricular tachycardia in patients with hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 1999 May;33(6):1596-601. [PubMed](#)

Epstein SE, Rosing DR. Verapamil: its potential for causing serious complications in patients with hypertrophic cardiomyopathy. *Circulation*. 1981 Sep;64(3):437-41. [PubMed](#)

Erwin JP 3rd, Nishimura RA, Lloyd MA, Tajik AJ. Dual chamber pacing for patients with hypertrophic obstructive cardiomyopathy: a clinical perspective in 2000. *Mayo Clin Proc*. 2000 Feb;75(2):173-80. [34 references] [PubMed](#)

Faber L, Seggewiss H, Welge D, Fassbender D, Schmidt HK, Gleichmann U, Horstkotte D. Echo-guided percutaneous septal ablation for symptomatic hypertrophic obstructive cardiomyopathy: 7 years of experience. *Eur J Echocardiogr*. 2004 Oct;5(5):347-55. [PubMed](#)

Fananapazir L, Chang AC, Epstein SE, McAreavey D. Prognostic determinants in hypertrophic cardiomyopathy. Prospective evaluation of a therapeutic strategy based on clinical, Holter, hemodynamic, and electrophysiological findings. *Circulation*. 1992 Sep;86(3):730-40. [PubMed](#)

Fernandes VL, Nielsen C, Nagueh SF, Herrin AE, Slifka C, Franklin J, Spencer WH 3rd. Follow-up of alcohol septal ablation for symptomatic hypertrophic obstructive cardiomyopathy the Baylor and Medical University of South Carolina experience 1996 to 2007. *JACC Cardiovasc Interv*. 2008 Oct;1(5):561-70. [PubMed](#)

Fifer MA, Vlahakes GJ. Management of symptoms in hypertrophic cardiomyopathy. *Circulation*. 2008 Jan 22;117(3):429-39. [118 references] [PubMed](#)

Firoozi S, Elliott PM, Sharma S, Murday A, Brecker SJ, Hamid MS, Sachdev B, Thaman R, McKenna WJ. Septal myotomy-myectomy and transcatheter septal alcohol ablation in hypertrophic obstructive cardiomyopathy. A comparison of clinical, haemodynamic and exercise outcomes. *Eur Heart J*. 2002 Oct;23(20):1617-24. [PubMed](#)

Fitzgerald-Butt SM, Byrne L, Gerhardt CA, Vannatta K, Hoffman TM, McBride KL. Parental knowledge and attitudes toward hypertrophic cardiomyopathy genetic testing. *Pediatr Cardiol*. 2010 Feb;31(2):195-202. [PubMed](#)

Flamm MD, Harrison DC, Hancock EW. Muscular subaortic stenosis. Prevention of outflow obstruction with propranolol. *Circulation*. 1968 Nov;38(5):846-58. [PubMed](#)

Fokstuen S, Lyle R, Munoz A, Gehrig C, Lerch R, Perrot A, Osterziel KJ, Geier C, Beghetti M, Mach F, Sztajzel J, Sigwart U, Antonarakis SE, Blouin JL. A DNA resequencing array for pathogenic mutation detection in hypertrophic cardiomyopathy. *Hum Mutat*. 2008 Jun;29(6):879-85. [PubMed](#)

Frank MJ, Abdulla AM, Canedo MI, Saylor RE. Long-term medical management of hypertrophic obstructive cardiomyopathy. *Am J Cardiol*. 1978 Dec;42(6):993-1001. [PubMed](#)

Frenneaux MP, Counihan PJ, Caforio AL, Chikamori T, McKenna WJ. Abnormal blood pressure response during exercise in hypertrophic cardiomyopathy. *Circulation*. 1990 Dec;82(6):1995-2002. [PubMed](#)

Fuster V, Ryden LE, Cannom DS, Crijns HJ, Curtis AB, Ellenbogen KA, Halperin JL, Kay GN, Le Huez JY, Lowe JE, Olsson SB, Prystowsky EN, Tamargo JL, Wann LS, Smith SC Jr, Priori SG, Estes NA 3rd, Ezekowitz MD, Jackman WM, January CT, Lowe JE, Page RL, Slotwimer DJ, Stevenson WG, Tracy CM, Jacobs AK, Anderson JL, Albert N, Buller CE, Creager MA, Ettinger SM, Guyton RA, Halperin JL, Hochman JS, Kushner FG, Ohman EM, Stevenson WG, Tarkington LG, Yancy CW, American College of Cardiology Foundation/American Heart Association Task Force. 2011 ACCF/AHA/HRS focused updates incorporated into the ACC/AHA/ESC 2006 guidelines for the management of patients with atrial fibrillation: a report of the American College of Cardiology Foundation/American Heart Association Task Force on [trunc]. *Circulation*. 2011 Mar 15;123(10):e269-367. [PubMed](#)

Gadler F, Linde C, Daubert C, McKenna W, Meisel E, Aliot E, Chojnowska L, Guize L, Gras D, Jeanrenaud X, Kappenberger L. Significant improvement of quality of life following atrioventricular synchronous pacing in patients with hypertrophic obstructive cardiomyopathy. Data from 1 year of follow-up. PIC study group. *Pacing In Cardiomyopathy. Eur Heart J*. 1999 Jul;20(14):1044-50. [PubMed](#)

Gaita F, Di Donna P, Olivetto I, Scaglione M, Ferrero I, Montefusco A, Caponi D, Conte MR, Nistri S, Cecchi F. Usefulness and safety of transcatheter ablation of atrial fibrillation in patients with hypertrophic cardiomyopathy. *Am J Cardiol*. 2007 Jun 1;99(11):1575-81. [PubMed](#)

Gajarski R, Naftel DC, Pahl E, Alejos J, Pearce FB, Kirklin JK, Zamberlan M, Dipchand AI, Pediatric Heart Transplant Study Investigators. Outcomes of pediatric patients with hypertrophic cardiomyopathy listed for transplant. *J Heart Lung Transplant*. 2009 Dec;28(12):1329-34. [PubMed](#)

Gange CA, Link MS, Maron MS. Utility of cardiovascular magnetic resonance in the diagnosis of Anderson-Fabry disease. *Circulation*. 2009 Sep 29;120(13):e96-7. [PubMed](#)

Gietzen FH, Leuner CJ, Obergassel L, Strunk-Mueller C, Kuhn H. Transcoronary ablation of septal hypertrophy for hypertrophic obstructive cardiomyopathy: feasibility, clinical benefit, and short term results in elderly patients. *Heart*. 2004 Jun;90(6):638-44. [PubMed](#)

Grigg LE, Wigle ED, Williams WG, Daniel LB, Rakowski H. Transesophageal Doppler echocardiography in obstructive hypertrophic cardiomyopathy: clarification of pathophysiology and importance in intraoperative decision making. *J Am Coll Cardiol*. 1992 Jul;20(1):42-52. [PubMed](#)

Haley JH, Sinak LJ, Tajik AJ, Ommen SR, Oh JK. Dynamic left ventricular outflow tract obstruction in acute coronary syndromes: an important cause of new systolic murmur and cardiogenic shock. *Mayo Clin Proc*. 1999 Sep;74(9):901-6. [PubMed](#)

Harris KM, Spirito P, Maron MS, Zenovich AG, Formisano F, Lesser JR, Mackey-Bojack S, Manning WJ, Udelson JE, Maron BJ. Prevalence, clinical profile, and significance of left ventricular remodeling in the end-stage phase of hypertrophic cardiomyopathy. *Circulation*. 2006 Jul 18;114(3):216-25. [PubMed](#)

Harrison DC, Braunwald DE, Glick G, Mason DT, Chidsey CA, Ross J Jr. Effects of Beta adrenergic blockade on the circulation with particular reference to observations in patients with hypertrophic subaortic stenosis. *Circulation*. 1964 Jan;29:84-98. [PubMed](#)

Hauser RG, Maron BJ, Marine JE, Lampert R, Kadish AH, Winters SL, Scher DL, Biria M, Kalia A. Safety and efficacy of transvenous high-voltage implantable cardioverter-defibrillator leads in high-risk hypertrophic cardiomyopathy patients. *Heart Rhythm*. 2008 Nov;5(11):1517-22. [PubMed](#)

Hershberger RE, Cowan J, Morales A, Siegfried JD. Progress with genetic cardiomyopathies: screening, counseling, and testing in dilated, hypertrophic, and arrhythmogenic right ventricular dysplasia/cardiomyopathy. *Circ Heart Fail*. 2009 May;2(3):253-61. [51 references] [PubMed](#)

Ho CY, Lever HM, DeSanctis R, Farver CF, Seidman JG, Seidman CE. Homozygous mutation in cardiac troponin T: implications for hypertrophic cardiomyopathy. *Circulation*. 2000 Oct 17;102(16):1950-5. [PubMed](#)

Ho CY, Sweitzer NK, McDonough B, Maron BJ, Casey SA, Seidman JG, Seidman CE, Solomon SD. Assessment of diastolic function with Doppler tissue imaging to predict genotype in preclinical hypertrophic cardiomyopathy. *Circulation*. 2002 Jun 25;105(25):2992-7. [PubMed](#)

Ingles J, Doolan A, Chiu C, Seidman J, Seidman C, Semsarian C. Compound and double mutations in patients with hypertrophic cardiomyopathy: implications for genetic testing and counselling. *J Med Genet*. 2005 Oct;42(10):e59. [PubMed](#)

Jassal DS, Neilan TG, Fifer MA, Palacios IF, Lowry PA, Vlahakes GJ, Picard MH, Yoerger DM. Sustained improvement in left ventricular diastolic function after alcohol septal ablation for hypertrophic obstructive cardiomyopathy. *Eur Heart J*. 2006 Aug;27(15):1805-10. [PubMed](#)

Jensen G, Uhrenholt A. Circulatory effects of intravenous disopyramide in heart failure. *J Int Med Res*. 1976;4(1 Suppl):42-5. [PubMed](#)

Jeschke B, Uhl K, Weist B, Schroder D, Meitinger T, Dohlemann C, Vosberg HP. A high risk phenotype of hypertrophic cardiomyopathy associated with a compound genotype of two mutated beta-myosin heavy chain genes. *Hum Genet*. 1998 Mar;102(3):299-304. [PubMed](#)

Kappenberger L, Linde C, Daubert C, McKenna W, Meisel E, Sadoul N, Chojnowska L, Guize L, Gras D, Jeanrenaud X, Ryden L. Pacing in hypertrophic obstructive cardiomyopathy. A randomized crossover study. PIC Study Group. *Eur Heart J*. 1997 Aug;18(8):1249-56. [PubMed](#)

Kilicaslan F, Verma A, Saad E, Themistoclakis S, Bonso A, Raviele A, Bozbas H, Andrews MW, Beheiry S, Hao S, Cummings JE, Marrouche NF, Lakkireddy D, Wazni O, Yamaji H, Saenz LC, Saliba W, Schweikert RA, Natale A. Efficacy of catheter ablation of atrial fibrillation in patients with hypertrophic obstructive cardiomyopathy. *Heart Rhythm*. 2006 Mar;3(3):275-80. [PubMed](#)

Kimball BP, Bui S, Wigle ED. Acute dose-response effects of intravenous disopyramide in hypertrophic obstructive cardiomyopathy. *Am Heart J*. 1993 Jun;125(6):1691-7. [PubMed](#)

Kleemann T, Becker T, Doenges K, Vater M, Senges J, Schneider S, Saggau W, Weisse U, Seidl K. Annual rate of transvenous defibrillation lead defects in implantable cardioverter-defibrillators over a period of >10 years. *Circulation*. 2007 May 15;115(19):2474-80. [PubMed](#)

Klues HG, Schiffrers A, Maron BJ. Phenotypic spectrum and patterns of left ventricular hypertrophy in hypertrophic cardiomyopathy: morphologic observations and significance as assessed by two-dimensional echocardiography in 600 patients. *J Am Coll Cardiol*. 1995 Dec;26(7):1699-708. [PubMed](#)

Krasnow N, Rolett E, Hood WB Jr, Yurchak PM, Gorlin R. Reversible obstruction of the ventricular outflow tract. *Am J Cardiol*. 1963 Jan;11:1-7. [PubMed](#)

Kuhn H, Lawrenz T, Lieder F, Leuner C, Strunk-Mueller C, Obergassel L, Bartelsmeier M, Stellbrink C. Survival after transcatheter ablation of septal hypertrophy in hypertrophic obstructive cardiomyopathy (TASH): a 10 year experience. *Clin Res Cardiol*. 2008 Apr;97(4):234-43. [PubMed](#)

Kwon DH, Kapadia SR, Tuzcu EM, Halley CM, Gorodeski EZ, Curtin RJ, Thamilarasan M, Smedira NG, Lytle BW, Lever HM, Desai MY. Long-term outcomes in high-risk symptomatic patients with hypertrophic cardiomyopathy undergoing alcohol septal ablation. *JACC Cardiovasc Interv*. 2008

Aug;1(4):432-8. [PubMed](#)

Lara M, Oakley GD, Rowbotham D. Potentially dangerous effect of disopyramide on atrioventricular conduction in a patient on digitalis. *Br Med J*. 1980 Jul 19;281(6234):198. [PubMed](#)

Maceira AM, Joshi J, Prasad SK, Moon JC, Perugini E, Harding I, Sheppard MN, Poole-Wilson PA, Hawkins PN, Pennell DJ. Cardiovascular magnetic resonance in cardiac amyloidosis. *Circulation*. 2005 Jan 18;111(2):186-93. [PubMed](#)

Maisel WH, Sweeney MO, Stevenson WG, Ellison KE, Epstein LM. Recalls and safety alerts involving pacemakers and implantable cardioverter-defibrillator generators. *JAMA*. 2001 Aug 15;286(7):793-9. [PubMed](#)

Maki S, Ikeda H, Muro A, Yoshida N, Shibata A, Koga Y, Inaizumi T. Predictors of sudden cardiac death in hypertrophic cardiomyopathy. *Am J Cardiol*. 1998 Sep 15;82(6):774-8. [PubMed](#)

Maron BJ, Ackerman MJ, Nishimura RA, Pyeritz RE, Towbin JA, Udelson JE. Task Force 4: HCM and other cardiomyopathies, mitral valve prolapse, myocarditis, and Marfan syndrome. *J Am Coll Cardiol*. 2005 Apr 19;45(8):1340-5. [49 references] [PubMed](#)

Maron BJ, Carney KP, Lever HM, Lewis JF, Barac I, Casey SA, Sherrid MV. Relationship of race to sudden cardiac death in competitive athletes with hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2003 Mar 19;41(6):974-80. [PubMed](#)

Maron BJ, Chaitman BR, Ackerman MJ, Bayes de Luna A, Corrado D, Crosson JE, Deal BJ, Driscoll DJ, Estes NA 3rd, Araujo CG, Liang DH, Mitten MJ, Myerburg RJ, Pelliccia A, Thompson PD, Towbin JA, Van Camp SP. Recommendations for physical activity and recreational sports participation for young patients with genetic cardiovascular diseases. *Circulation*. 2004 Jun 8;109(22):2807-16. [86 references] [PubMed](#)

Maron BJ, Doerer JJ, Haas TS, Tierney DM, Mueller FO. Sudden deaths in young competitive athletes: analysis of 1866 deaths in the United States, 1980-2006. *Circulation*. 2009 Mar 3;119(8):1085-92. [PubMed](#)

Maron BJ, Epstein SE, Roberts WC. Causes of sudden death in competitive athletes. *J Am Coll Cardiol*. 1986 Jan;7(1):204-14. [72 references] [PubMed](#)

Maron BJ, McKenna WJ, Danielson GK, Kappenberger LJ, Kuhn HJ, Seidman CE, Shah PM, Spencer WH 3rd, Spirito P, Ten Cate FJ, Wigle ED, Task Force on Clinical Expert Consensus Documents. American College of Cardiology, Committee for Practice Guidelines. European Society of Cardiology. American College of Cardiology/European Society of Cardiology clinical expert consensus document on hypertrophic cardiomyopathy. A report of the American College of Cardiology Foundation Task Force on Clinical Expert Consensus Documents [trunc]. *J Am Coll Cardiol*. 2003 Nov 5;42(9):1687-713. [273 references] [PubMed](#)

Maron BJ, Niiimura H, Casey SA, Soper MK, Wright GB, Seidman JG, Seidman CE. Development of left ventricular hypertrophy in adults in hypertrophic cardiomyopathy caused by cardiac myosin-binding protein C gene mutations. *J Am Coll Cardiol*. 2001 Aug;38(2):315-21. [PubMed](#)

Maron BJ, Nishimura RA, McKenna WJ, Rakowski H, Josephson ME, Kieval RS. Assessment of permanent dual-chamber pacing as a treatment for drug-refractory symptomatic patients with obstructive hypertrophic cardiomyopathy. A randomized, double-blind, crossover study (M-PATHY). *Circulation*. 1999 Jun 8;99(22):2927-33. [PubMed](#)

Maron BJ, Olivotto I, Bellone P, Conte MR, Cecchi F, Flygenring BP, Casey SA, Gohman TE, Bongioanni S, Spirito P. Clinical profile of stroke in 900 patients with hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2002 Jan 16;39(2):301-7. [PubMed](#)

Maron BJ, Olivotto I, Spirito P, Casey SA, Bellone P, Gohman TE, Graham KJ, Burton DA, Cecchi F. Epidemiology of hypertrophic cardiomyopathy-related death: revisited in a large non-referral-based patient population. *Circulation*. 2000 Aug 22;102(8):858-64. [PubMed](#)

Maron BJ, Savage DD, Wolfson JK, Epstein SE. Prognostic significance of 24 hour ambulatory electrocardiographic monitoring in patients with hypertrophic cardiomyopathy: a prospective study. *Am J Cardiol.* 1981 Aug;48(2):252-7. [PubMed](#)

Maron BJ, Seidman JG, Seidman CE. Proposal for contemporary screening strategies in families with hypertrophic cardiomyopathy. *J Am Coll Cardiol.* 2004 Dec 7;44(11):2125-32. [92 references] [PubMed](#)

Maron BJ, Shen WK, Link MS, Epstein AE, Almquist AK, Daubert JP, Bardy GH, Favale S, Rea RF, Boriani G, Estes NA 3rd, Spirito P. Efficacy of implantable cardioverter-defibrillators for the prevention of sudden death in patients with hypertrophic cardiomyopathy. *N Engl J Med.* 2000 Feb 10;342(6):365-73. [PubMed](#)

Maron BJ, Spirito P, Shen WK, Haas TS, Formisano F, Link MS, Epstein AE, Almquist AK, Daubert JP, Lawrenz T, Boriani G, Estes NA 3rd, Favale S, Piccininno M, Winters SL, Santini M, Betocchi S, Arribas F, Sherrid MV, Buja G, Sensarian C, Bruzzi P. Implantable cardioverter-defibrillators and prevention of sudden cardiac death in hypertrophic cardiomyopathy. *JAMA.* 2007 Jul 25;298(4):405-12. [PubMed](#)

Maron BJ, Spirito P. Implantable defibrillators and prevention of sudden death in hypertrophic cardiomyopathy. *J Cardiovasc Electrophysiol.* 2008 Oct;19(10):1118-26. [89 references] [PubMed](#)

Maron BJ, Spirito P. Implications of left ventricular remodeling in hypertrophic cardiomyopathy. *Am J Cardiol.* 1998 Jun 1;81(11):1339-44. [78 references] [PubMed](#)

Maron BJ. Contemporary insights and strategies for risk stratification and prevention of sudden death in hypertrophic cardiomyopathy. *Circulation.* 2010 Jan 26;121(3):445-56. [90 references] [PubMed](#)

Maron BJ. Hypertrophic cardiomyopathy: a systematic review. *JAMA.* 2002 Mar 13;287(10):1308-20. [207 references] [PubMed](#)

Maron BJ. Risk stratification and role of implantable defibrillators for prevention of sudden death in patients with hypertrophic cardiomyopathy. *Circ J.* 2010 Nov;74(11):2271-82. [PubMed](#)

Maron BJ. Sudden death in young athletes. *N Engl J Med.* 2003 Sep 11;349(11):1064-75. [100 references] [PubMed](#)

Maron MS, Appelbaum E, Harrigan CJ, Buros J, Gibson CM, Hanna C, Lesser JR, Udelson JE, Manning WJ, Maron BJ. Clinical profile and significance of delayed enhancement in hypertrophic cardiomyopathy. *Circ Heart Fail.* 2008 Sep;1(3):184-91. [PubMed](#)

Maron MS, Finley JJ, Bos JM, Hauser TH, Manning WJ, Haas TS, Lesser JR, Udelson JE, Ackerman MJ, Maron BJ. Prevalence, clinical significance, and natural history of left ventricular apical aneurysms in hypertrophic cardiomyopathy. *Circulation.* 2008 Oct 7;118(15):1541-9. [PubMed](#)

Maron MS, Lesser JR, Maron BJ. Management implications of massive left ventricular hypertrophy in hypertrophic cardiomyopathy significantly underestimated by echocardiography but identified by cardiovascular magnetic resonance. *Am J Cardiol.* 2010 Jun 15;105(12):1842-3. [PubMed](#)

Maron MS, Maron BJ, Harrigan C, Buros J, Gibson CM, Olivotto I, Biller L, Lesser JR, Udelson JE, Manning WJ, Appelbaum E. Hypertrophic cardiomyopathy phenotype revisited after 50 years with cardiovascular magnetic resonance. *J Am Coll Cardiol.* 2009 Jul 14;54(3):220-8. [PubMed](#)

Maron MS, Olivotto I, Betocchi S, Casey SA, Lesser JR, Losi MA, Cecchi F, Maron BJ. Effect of left ventricular outflow tract obstruction on clinical outcome in hypertrophic cardiomyopathy. *N Engl J Med.* 2003 Jan 23;348(4):295-303. [PubMed](#)

Maron MS, Olivotto I, Zenovich AG, Link MS, Pandian NG, Kuvin JT, Nistri S, Cecchi F, Udelson JE, Maron BJ. Hypertrophic cardiomyopathy is predominantly a disease of left ventricular outflow tract obstruction. *Circulation.* 2006 Nov 21;114(21):2232-9. [PubMed](#)

Marwick TH, Stewart WJ, Lever HM, Lytle BW, Rosenkranz ER, Duffy CI, Salcedo EE. Benefits of intraoperative echocardiography in the surgical management of hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 1992 Nov 1;20(5):1066-72. [PubMed](#)

McKenna W, Deanfield J, Faruqi A, England D, Oakley C, Goodwin J. Prognosis in hypertrophic cardiomyopathy: role of age and clinical, electrocardiographic and hemodynamic features. *Am J Cardiol*. 1981 Mar;47(3):532-8. [PubMed](#)

Michels M, Soliman OI, Pfeifferkorn J, Hoedemackers YM, Kofflard MJ, Dooijes D, Majoor-Krakauer D, Ten Cate FJ. Disease penetrance and risk stratification for sudden cardiac death in asymptomatic hypertrophic cardiomyopathy mutation carriers. *Eur Heart J*. 2009 Nov;30(21):2593-8. [PubMed](#)

Michie S, Allanson A, Armstrong D, Weinman J, Bobrow M, Marteau TM. Objectives of genetic counselling: differing views of purchasers, providers and users. *J Public Health Med*. 1998 Dec;20(4):404-8. [PubMed](#)

Michie S, French D, Allanson A, Bobrow M, Marteau TM. Information recall in genetic counselling: a pilot study of its assessment. *Patient Educ Couns*. 1997 Sep-Oct;32(1-2):93-100. [PubMed](#)

Monakier D, Woo A, Puri T, Schwartz L, Ross J, Jamorski M, Yang H, Liu Z, Vannan M, Wigle ED, Rakowski H. Usefulness of myocardial contrast echocardiographic quantification of risk area for predicting postprocedural complications in patients undergoing septal ethanol ablation for obstructive hypertrophic cardiomyopathy. *Am J Cardiol*. 2004 Dec 15;94(12):1515-22. [PubMed](#)

Montserrat L, Elliott PM, Gimeno JR, Sharma S, Penas-Lado M, McKenna WJ. Non-sustained ventricular tachycardia in hypertrophic cardiomyopathy: an independent marker of sudden death risk in young patients. *J Am Coll Cardiol*. 2003 Sep 3;42(5):873-9. [PubMed](#)

Moolman JC, Corfield VA, Posen B, Ngumbela K, Seidman C, Brink PA, Watkins H. Sudden death due to troponin T mutations. *J Am Coll Cardiol*. 1997 Mar 1;29(3):549-55. [PubMed](#)

Moon JC, Fisher NG, McKenna WJ, Pennell DJ. Detection of apical hypertrophic cardiomyopathy by cardiovascular magnetic resonance in patients with non-diagnostic echocardiography. *Heart*. 2004 Jun;90(6):645-9. [PubMed](#)

Moon JC, McKenna WJ, McCrohon JA, Elliott PM, Smith GC, Pennell DJ. Toward clinical risk assessment in hypertrophic cardiomyopathy with gadolinium cardiovascular magnetic resonance. *J Am Coll Cardiol*. 2003 May 7;41(9):1561-7. [PubMed](#)

Moon JC, Sachdev B, Elkington AG, McKenna WJ, Mehta A, Pennell DJ, Leed PJ, Elliott PM. Gadolinium enhanced cardiovascular magnetic resonance in Anderson-Fabry disease. Evidence for a disease specific abnormality of the myocardial interstitium. *Eur Heart J*. 2003 Dec;24(23):2151-5. [PubMed](#)

Morady F, Scheinman MM, Desai J. Disopyramide. *Ann Intern Med*. 1982 Mar;96(3):337-43. [71 references] [PubMed](#)

Morita H, Rehm HL, Menesses A, McDonough B, Roberts AE, Kucherlapati R, Towbin JA, Seidman JG, Seidman CE. Shared genetic causes of cardiac hypertrophy in children and adults. *N Engl J Med*. 2008 May 1;358(18):1899-908. [PubMed](#)

Nagueh SF, Lakkis NM, He ZX, Middleton KJ, Killip D, Zoghbi WA, Quinones MA, Roberts R, Verani MS, Kleiman NS, Spencer WH 3d. Role of myocardial contrast echocardiography during nonsurgical septal reduction therapy for hypertrophic obstructive cardiomyopathy. *J Am Coll Cardiol*. 1998 Jul;32(1):225-9. [PubMed](#)

Nagueh SF, Ommen SR, Lakkis NM, Killip D, Zoghbi WA, Schaff HV, Danielson GK, Quinones MA, Tajik AJ, Spencer WH. Comparison of ethanol septal reduction therapy with surgical myectomy for the treatment of hypertrophic obstructive cardiomyopathy. *J Am Coll Cardiol*. 2001 Nov 15;38(6):1701-6. [PubMed](#)

Niimura H, Bachinski LL, Sangwatanaroj S, Watkins H, Chudley AE, McKenna W, Kristinsson A, Roberts R, Sole M, Maron BJ, Seidman JG, Seidman CE. Mutations in the gene for cardiac myosin-binding protein C and late-onset familial hypertrophic cardiomyopathy. *N Engl J Med*. 1998 Apr 30;338(18):1248-57. [PubMed](#)

Nishimura RA, Trusty JM, Hayes DL, Ilstrup DM, Larson DR, Hayes SN, Allison TG, Tajik AJ. Dual-chamber pacing for hypertrophic cardiomyopathy: a randomized, double-blind, crossover trial. *J Am Coll Cardiol*. 1997 Feb;29(2):435-41. [PubMed](#)

Offit K, Groeger E, Turner S, Wadsworth EA, Weiser MA. The "duty to warn" a patient's family members about hereditary disease risks. *JAMA*. 2004 Sep 22;292(12):1469-73. [PubMed](#)

O'Hanlon R, Grasso A, Roughton M, Moon JC, Clark S, Wage R, Webb J, Kulkarni M, Dawson D, Sulaibeekh L, Chandrasekaran B, Bucciarelli-Ducci C, Pasquale F, Cowie MR, McKenna WJ, Sheppard MN, Elliott PM, Pennell DJ, Prasad SK. Prognostic significance of myocardial fibrosis in hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2010 Sep 7;56(11):867-74. [PubMed](#)

Olivotto I, Cecchi F, Casey SA, Dolara A, Traverse JH, Maron BJ. Impact of atrial fibrillation on the clinical course of hypertrophic cardiomyopathy. *Circulation*. 2001 Nov 20;104(21):2517-24. [PubMed](#)

Olivotto I, Girolami F, Ackerman MJ, Nistri S, Bos JM, Zachara E, Ommen SR, Theis JL, Vaubel RA, Re F, Armentano C, Poggesi C, Torricelli F, Cecchi F. Myofilament protein gene mutation screening and outcome of patients with hypertrophic cardiomyopathy. *Mayo Clin Proc*. 2008 Jun;83(6):630-8. [PubMed](#)

Olivotto I, Maron BJ, Monterege A, Mazzuoli F, Dolara A, Cecchi F. Prognostic value of systemic blood pressure response during exercise in a community-based patient population with hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 1999 Jun;33(7):2044-51. [PubMed](#)

Ommen SR, Maron BJ, Olivotto I, Maron MS, Cecchi F, Betocchi S, Gersh BJ, Ackerman MJ, McCully RB, Dearani JA, Schaff HV, Danielson GK, Tajik AJ, Nishimura RA. Long-term effects of surgical septal myectomy on survival in patients with obstructive hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2005 Aug 2;46(3):470-6. [PubMed](#)

Ommen SR, Nishimura RA, Squires RW, Schaff HV, Danielson GK, Tajik AJ. Comparison of dual-chamber pacing versus septal myectomy for the treatment of patients with hypertrophic obstructive cardiomyopathy: a comparison of objective hemodynamic and exercise end points. *J Am Coll Cardiol*. 1999 Jul;34(1):191-6. [PubMed](#)

Ommen SR, Shah PM, Tajik AJ. Left ventricular outflow tract obstruction in hypertrophic cardiomyopathy: past, present and future. *Heart*. 2008 Oct;94(10):1276-81. [74 references] [PubMed](#)

Pelliccia A, Fagard R, Bjornstad HH, Anastassakis A, Arbustini E, Assanelli D, Biffi A, Borjesson M, Carre F, Corrado D, Delise P, Dorwarth U, Hirth A, Heidbuchel H, Hoffmann E, Mellwig KP, Panhuyzen-Goedkoop N, Pisani A, Solberg EE, van-Buuren F, Vanhees L, Blomstrom-Lundqvist C, Deligiannis A, Dugmore D, Glikson M, Hoff PI, Hoffmann A, Hoffmann E, Horstkotte D, Nordrehaug JE, Oudhof J, McKenna WJ, Penco M, Priori S, Reybrouck T, Senden J, Spataro A, Thiene G, Study Group of Sports Cardiology of the Working Group of Cardiac Rehabilitation, Working Group of Myocardial and Pericardial Diseases of the European Society of. Recommendations for competitive sports participation in athletes with cardiovascular disease: a consensus document from the Study Group of Sports Cardiology of the Working Group of Cardiac Rehabilitation and Exercise Physiology [trunc] . *Eur Heart J*. 2005 Jul;26(14):1422-45. [98 references] [PubMed](#)

Pierce GE, Morrow AG, Braunwald E. Idiopathic hypertrophic subaortic stenosis. 3. Intraoperative studies of the mechanism of obstruction and its hemodynamic consequences. *Circulation*. 1964 Nov;30(Suppl 4):152. [PubMed](#)

Pollick C, Kimball B, Henderson M, Wigle ED. Disopyramide in hypertrophic cardiomyopathy. I. Hemodynamic assessment after intravenous administration. *Am J Cardiol*. 1988 Dec 1;62(17):1248-51. [PubMed](#)

Pollick C. Disopyramide in hypertrophic cardiomyopathy. II. Noninvasive assessment after oral administration. *Am J Cardiol.* 1988 Dec 1;62(17):1252-5. [PubMed](#)

Qin JX, Shiota T, Lever HM, Kapadia SR, Sitges M, Rubin DN, Bauer F, Greenberg NL, Agler DA, Drinko JK, Martin M, Tuzcu EM, Smedira NG, Lytle B, Thomas JD. Outcome of patients with hypertrophic obstructive cardiomyopathy after percutaneous transluminal septal myocardial ablation and septal myectomy surgery. *J Am Coll Cardiol.* 2001 Dec;38(7):1994-2000. [PubMed](#)

Ralph-Edwards A, Woo A, McCrindle BW, Shapero JL, Schwartz L, Rakowski H, Wigle ED, Williams WG. Hypertrophic obstructive cardiomyopathy: comparison of outcomes after myectomy or alcohol ablation adjusted by propensity score. *J Thorac Cardiovasc Surg.* 2005 Feb;129(2):351-8. [PubMed](#)

Redberg RF, Benjamin EJ, Bittner V, Braun LT, Goff DC Jr, Havas S, Labarthe DR, Limacher MC, Lloyd-Jones DM, Mora S, Pearson TA, Radford MJ, Smetana GW, Spertus JA, Swegler EW, American Academy of Family Physicians, American Association of Cardiovascular and Pulmonary Rehabilitation, Preventive Cardiovascular Nurses Association. ACCF/AHA 2009 performance measures for primary prevention of cardiovascular disease in adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Performance Measures [trunc]. *J Am Coll Cardiol.* 2009 Sep 29;54(14):1364-405. [PubMed](#)

Rickers C, Wilke NM, Jerosch-Herold M, Casey SA, Panse P, Panse N, Weil J, Zenovich AG, Maron BJ. Utility of cardiac magnetic resonance imaging in the diagnosis of hypertrophic cardiomyopathy. *Circulation.* 2005 Aug 9;112(6):855-61. [PubMed](#)

Robertson CE, Miller HC. Extreme tachycardia complicating the use of disopyramide in atrial flutter. *Br Heart J.* 1980 Nov;44(5):602-3. [PubMed](#)

Rosenzweig A, Watkins H, Hwang DS, Miri M, McKenna W, Traill TA, Seidman JG, Seidman CE. Preclinical diagnosis of familial hypertrophic cardiomyopathy by genetic analysis of blood lymphocytes. *N Engl J Med.* 1991 Dec 19;325(25):1753-60. [PubMed](#)

Rosing DR, Condit JR, Maron BJ, Kent KM, Leon MB, Bonow RO, Lipson LC, Epstein SE. Verapamil therapy: a new approach to the pharmacologic treatment of hypertrophic cardiomyopathy: III. Effects of long-term administration. *Am J Cardiol.* 1981 Sep;48(3):545-53. [PubMed](#)

Rosing DR, Kent KM, Borer JS, Seides SF, Maron BJ, Epstein SE. Verapamil therapy: a new approach to the pharmacologic treatment of hypertrophic cardiomyopathy. I. Hemodynamic effects. *Circulation.* 1979 Dec;60(6):1201-7. [PubMed](#)

Rosing DR, Kent KM, Maron BJ, Epstein SE. Verapamil therapy: a new approach to the pharmacologic treatment of hypertrophic cardiomyopathy. II. Effects on exercise capacity and symptomatic status. *Circulation.* 1979 Dec;60(6):1208-13. [PubMed](#)

Rubinshtein R, Glockner JF, Ommen SR, Araoz PA, Ackerman MJ, Sorajja P, Bos JM, Tajik AJ, Valeti US, Nishimura RA, Gersh BJ. Characteristics and clinical significance of late gadolinium enhancement by contrast-enhanced magnetic resonance imaging in patients with hypertrophic cardiomyopathy. *Circ Heart Fail.* 2010 Jan;3(1):51-8. [PubMed](#)

Sadoul N, Prasad K, Elliott PM, Bannerjee S, Frenneaux MP, McKenna WJ. Prospective prognostic assessment of blood pressure response during exercise in patients with hypertrophic cardiomyopathy. *Circulation.* 1997 Nov 4;96(9):2987-91. [PubMed](#)

Schwartz ML, Cox GF, Lin AE, Korson MS, Perez-Atayde A, Lacro RV, Lipshultz SE. Clinical approach to genetic cardiomyopathy in children. *Circulation.* 1996 Oct 15;94(8):2021-38. [214 references] [PubMed](#)

Sherrid M, Delia E, Dwyer E. Oral disopyramide therapy for obstructive hypertrophic cardiomyopathy. *Am J Cardiol.* 1988 Nov 15;62(16):1085-8. [PubMed](#)

Sherrid MV, Barac I, McKenna WJ, Elliott PM, Dickie S, Chojnowska L, Casey S, Maron BJ. Multicenter study of the efficacy and safety of disopyramide in obstructive hypertrophic cardiomyopathy. *J Am Coll Cardiol.* 2005 Apr 19;45(8):1251-8. [PubMed](#)

Slade AK, Sadoul N, Shapiro L, Chojnowska L, Simon JP, Saumarez RC, Dodinot B, Camm AJ, McKenna WJ, Aliot E. DDD pacing in hypertrophic cardiomyopathy: a multicentre clinical experience. *Heart*. 1996 Jan;75(1):44-9. [PubMed](#)

Smedira NG, Lytle BW, Lever HM, Rajeswaran J, Krishnaswamy G, Kaple RK, Dolney DO, Blackstone EH. Current effectiveness and risks of isolated septal myectomy for hypertrophic obstructive cardiomyopathy. *Ann Thorac Surg*. 2008 Jan;85(1):127-33. [PubMed](#)

Sonnenblick EH, Williams JF Jr, Glick G, Mason DT, Braunwald E. Studies on digitalis. XV. Effects of cardiac glycosides on myocardial force-velocity relations in the nonfailing human heart. *Circulation*. 1966 Sep;34(3):532-9. [PubMed](#)

Soor GS, Luk A, Ahn E, Abraham JR, Woo A, Ralph-Edwards A, Butany J. Hypertrophic cardiomyopathy: current understanding and treatment objectives. *J Clin Pathol*. 2009 Mar;62(3):226-35. [85 references] [PubMed](#)

Sorajja P, Nishimura RA, Gersh BJ, Dearani JA, Hodge DO, Wiste HJ, Ommen SR. Outcome of mildly symptomatic or asymptomatic obstructive hypertrophic cardiomyopathy: a long-term follow-up study. *J Am Coll Cardiol*. 2009 Jul 14;54(3):234-41. [PubMed](#)

Sorajja P, Nishimura RA, Ommen SR, Ackerman MJ, Tajik AJ, Gersh BJ. Use of echocardiography in patients with hypertrophic cardiomyopathy: clinical implications of massive hypertrophy. *J Am Soc Echocardiogr*. 2006 Jun;19(6):788-95. [PubMed](#)

Sorajja P, Valeti U, Nishimura RA, Ommen SR, Rihal CS, Gersh BJ, Hodge DO, Schaff HV, Holmes DR Jr. Outcome of alcohol septal ablation for obstructive hypertrophic cardiomyopathy. *Circulation*. 2008 Jul 8;118(2):131-9. [PubMed](#)

Spada M, Pagliardini S, Yasuda M, Tükel T, Thiagarajan G, Sakuraba H, Ponzone A, Desnick RJ. High incidence of later-onset fabry disease revealed by newborn screening. *Am J Hum Genet*. 2006 Jul;79(1):31-40. [PubMed](#)

Spirito P, Autore C, Rapezzi C, Bernabo P, Badagliacca R, Maron MS, Bongioanni S, Coccio F, Estes NA, Barilla CS, Biagini E, Quarta G, Conte MR, Bruzzi P, Maron BJ. Syncope and risk of sudden death in hypertrophic cardiomyopathy. *Circulation*. 2009 Apr 7;119(13):1703-10. [PubMed](#)

Spirito P, Bellone P, Harris KM, Bernabo P, Bruzzi P, Maron BJ. Magnitude of left ventricular hypertrophy and risk of sudden death in hypertrophic cardiomyopathy. *N Engl J Med*. 2000 Jun 15;342(24):1778-85. [PubMed](#)

Spirito P, Seidman CE, McKenna WJ, Maron BJ. The management of hypertrophic cardiomyopathy. *N Engl J Med*. 1997 Mar 13;336(11):775-85. [PubMed](#)

Stenson RE, Flamm MD Jr, Harrison DC, Hancock EW. Hypertrophic subaortic stenosis. Clinical and hemodynamic effects of long-term propranolol therapy. *Am J Cardiol*. 1973 Jun;31(6):763-73. [PubMed](#)

Swanton RH, Brooksby IA, Jenkins BS, Webb-Peploe MM. Hemodynamic studies of beta blockade in hypertrophic obstructive cardiomyopathy. *Eur J Cardiol*. 1977 Jun;5(4):327-41. [PubMed](#)

Tendera M, Wycisk A, Schneeweiss A, Polonski L, Wodniecki J. Effect of sotalol on arrhythmias and exercise tolerance in patients with hypertrophic cardiomyopathy. *Cardiology*. 1993;82(5):335-42. [PubMed](#)

Theodoro DA, Danielson GK, Feldt RH, Anderson BJ. Hypertrophic obstructive cardiomyopathy in pediatric patients: results of surgical treatment. *J Thorac Cardiovasc Surg*. 1996 Dec;112(6):1589-97; discussion 1597-. [PubMed](#)

Towbin JA. Cardiomyopathy and heart transplantation in children. *Curr Opin Cardiol*. 2002 May;17(3):274-9. [69 references] [PubMed](#)

van der Lee C, Scholzel B, ten Berg JM, Geleijnse ML, Idzerda HH, van Domburg RT, Vletter WB, Serruys PW, ten Cate FJ. Usefulness of clinical, echocardiographic, and procedural characteristics to predict outcome after percutaneous transluminal septal myocardial ablation. *Am J Cardiol*. 2008 May 1;101(9):1315-20. [PubMed](#)

Van Driest SL, Ackerman MJ, Ommen SR, Shakur R, Will ML, Nishimura RA, Tajik AJ, Gersh BJ. Prevalence and severity of "benign" mutations in the beta-myosin heavy chain, cardiac troponin T, and alpha-tropomyosin genes in hypertrophic cardiomyopathy. *Circulation*. 2002 Dec 10;106(24):3085-90. [PubMed](#)

Van Driest SL, Jaeger MA, Ommen SR, Will ML, Gersh BJ, Tajik AJ, Ackerman MJ. Comprehensive analysis of the beta-myosin heavy chain gene in 389 unrelated patients with hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2004 Aug 4;44(3):602-10. [PubMed](#)

Van Driest SL, Vasile VC, Ommen SR, Will ML, Tajik AJ, Gersh BJ, Ackerman MJ. Myosin binding protein C mutations and compound heterozygosity in hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2004 Nov 2;44(9):1903-10. [PubMed](#)

Whalen RE, Cohen AI, Sumner RG, McIntosh HD. Demonstration of the dynamic nature of idiopathic hypertrophic subaortic stenosis. *Am J Cardiol*. 1963 Jan;11:8-17. [PubMed](#)

Wigle ED, Adelman AG, Felderhof CH. Medical and surgical treatment of the cardiomyopathies. *Circ Res*. 1974 Aug;35(2):suppl II:196-207. [PubMed](#)

Wigle ED, David PR, Labrosse CJ, McMeekan J. Muscular subaortic stenosis; the interrelation of wall tension, outflow tract "distending pressure" and orifice radius. *Am J Cardiol*. 1965 Jun;15:761-72. [PubMed](#)

Wigle ED, Rakowski H, Kimball BP, Williams WG. Hypertrophic cardiomyopathy. Clinical spectrum and treatment. *Circulation*. 1995 Oct 1;92(7):1680-92. [209 references] [PubMed](#)

Wigle ED, Sasson Z, Henderson MA, Ruddy TD, Fulop J, Rakowski H, Williams WG. Hypertrophic cardiomyopathy. The importance of the site and the extent of hypertrophy. A review. *Prog Cardiovasc Dis*. 1985 Jul-Aug;28(1):1-83. [271 references] [PubMed](#)

Woo A, Rakowski H, Liew JC, Zhao MS, Liew CC, Parker TG, Zeller M, Wigle ED, Sole MJ. Mutations of the beta myosin heavy chain gene in hypertrophic cardiomyopathy: critical functional sites determine prognosis. *Heart*. 2003 Oct;89(10):1179-85. [PubMed](#)

Woo A, Williams WG, Choi R, Wigle ED, Rozenblyum E, Fedwick K, Siu S, Ralph-Edwards A, Rakowski H. Clinical and echocardiographic determinants of long-term survival after surgical myectomy in obstructive hypertrophic cardiomyopathy. *Circulation*. 2005 Apr 26;111(16):2033-41. [PubMed](#)

Yoerger DM, Picard MH, Palacios IF, Vlahakes GJ, Lowry PA, Fifer MA. Time course of pressure gradient response after first alcohol septal ablation for obstructive hypertrophic cardiomyopathy. *Am J Cardiol*. 2006 May 15;97(10):1511-4. [PubMed](#)

Yu EH, Omran AS, Wigle ED, Williams WG, Siu SC, Rakowski H. Mitral regurgitation in hypertrophic obstructive cardiomyopathy: relationship to obstruction and relief with myectomy. *J Am Coll Cardiol*. 2000 Dec;36(7):2219-25. [PubMed](#)

Type of Evidence Supporting the Recommendations

The type of supporting evidence is identified and graded for each recommendation (see the "Major Recommendations" field).

The recommendations listed in the original guideline document are, whenever possible, evidence based.

Benefits/Harms of Implementing the Guideline Recommendations

Potential Benefits

Appropriate and effective assessment and treatment of hypertrophic cardiomyopathy

Potential Harms

- Beta-blocking drugs should be used with caution in patients with sinus bradycardia or severe conduction disease.
- Verapamil should be used with caution in patients with high gradients, advanced heart failure, or sinus bradycardia.
- Angiotensin-converting enzyme inhibitors or angiotensin receptor blockers should be used cautiously (if at all) in patients with resting or provokable LVOT obstruction.
- Administration of beta-blocking drugs with either verapamil or diltiazem should also be performed with caution because of the potential for high-grade atrioventricular block.
- Class IC antiarrhythmic agents were associated with an increased mortality in patients with coronary artery disease. Thus, caution is advised when these agents are prescribed for patients with hypertrophic cardiomyopathy (HCM) and their use should probably be limited to individuals with an implantable cardioverter-defibrillator (ICD).
- Disopyramide may cause anticholinergic side effects that can be managed with dose reduction.
- Complications following myectomy are rare when performed in experienced centers. The risk of complete heart block is approximately 2% with myectomy (higher in patients with preexisting right bundle-branch block), but in myectomy patients who have had previous alcohol septal ablation, risk is much higher (50% to 85%). Iatrogenic ventricular septal defect occurs in <1% of patients. Finally, the risk of aortic valve or mitral valve injury is also low (<1%), particularly when myectomy is performed by an experienced operator.
- In approximately half of patients undergoing alcohol septal ablation, temporary complete atrioventricular block occurs during the procedure. Persistent complete heart block prompting implantation of a permanent pacemaker occurs in 10% to 20% of patients based on the available data. Approximately 5% of patients have sustained ventricular tachyarrhythmias during hospitalization. The in-hospital mortality rate is up to 2%. Because of the potential for creating a ventricular septal defect, septal ablation should not be performed if the target septal thickness is ≤ 15 mm.
- Potential early problems of ICD therapy in HCM may include pneumothorax, pericardial effusion, pocket hematoma, acute pocket infection, and/or lead dislodgment. Late complications include upper extremity deep venous thrombosis, lead dislodgment, infection, high defibrillation threshold necessitating lead revision, and inappropriate shocks, that is, shocks triggered by supraventricular arrhythmias, sinus tachycardia, lead fractures or dislodgment, oversensing, double counting, and programming malfunctions.

Contraindications

Contraindications

- Dihydropyridine class calcium channel blockers (e.g., nifedipine) should not be used in patients with obstructive physiology because their vasodilatory effects may aggravate outflow obstruction.
- Dabigatran should not be used in patients with prosthetic valves, hemodynamically significant valve disease, advanced liver failure, or severe renal failure (creatinine clearance <15 mL/min).

Qualifying Statements

Qualifying Statements

- The American College of Cardiology Foundation/American Heart Association (ACCF/AHA) practice guidelines are intended to assist healthcare providers in clinical decision making by describing a range of generally acceptable approaches for the diagnosis, management, and prevention of specific diseases or conditions. These practice guidelines represent a consensus of expert opinion after a thorough review of the available current scientific evidence and are intended to improve patient care. The guidelines attempt to define practices that meet the needs of most patients in most circumstances. The ultimate judgment regarding care of a particular patient must be made by the healthcare provider and patient in light of all the circumstances presented by that patient. Thus, there are situations in which deviations from these guidelines may be appropriate. Clinical decision making should consider the quality and availability of expertise in the area where care is provided. When these guidelines are used as the basis for regulatory or payer decisions, the goal should be improvement in quality of care. The Task Force recognizes that situations arise for which additional data are needed to better inform patient care; these areas will be identified within each respective guideline when appropriate.
- Prescribed courses of treatment in accordance with these recommendations are effective only if they are followed. Because lack of patient understanding and adherence may adversely affect outcomes, physicians and other healthcare providers should make every effort to engage the patient's active participation in prescribed medical regimens and lifestyles.

Implementation of the Guideline

Description of Implementation Strategy

An implementation strategy was not provided.

Implementation Tools

Clinical Algorithm

Pocket Guide/Reference Cards

Quick Reference Guides/Physician Guides

Slide Presentation

For information about availability, see the *Availability of Companion Documents* and *Patient Resources* fields below.

Institute of Medicine (IOM) National Healthcare Quality Report Categories

IOM Care Need

Living with Illness

Staying Healthy

IOM Domain

Effectiveness

Patient-centeredness

Identifying Information and Availability

Bibliographic Source(s)

Gersh BJ, Maron BJ, Bonow RO, Dearani JA, Fifer MA, Link MS, Naidu SS, Nishimura RA, Ommen SR, Rakowski H, Seidman CE, Towbin JA, Udelson JE, Yancy CW. 2011 ACCF/AHA guideline for the diagnosis and treatment of hypertrophic cardiomyopathy: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol. 2011 Dec 13;58(25):e212-60. [453 references] [PubMed](#)

Adaptation

Not applicable: The guideline was not adapted from another source.

Date Released

2011 Dec 13

Guideline Developer(s)

American College of Cardiology Foundation - Medical Specialty Society

Source(s) of Funding

The work of the writing committee was supported exclusively by the American College of Cardiology Foundation (ACCF) and American Heart Association (AHA) without commercial support.

Guideline Committee

American College of Cardiology Foundation/American Heart Association (ACCF/AHA) Task Force on Practice Guidelines

Composition of Group That Authored the Guideline

Writing Committee Members: Bernard J. Gersh, MB, CHB, DPhil, FACC, FAHA (*Co-Chair*); Barry J. Maron, MD, FACC (*Co-Chair*); Robert O. Bonow, MD, MACC, FAHA; Joseph A. Dearani, MD, FACC; Michael A. Fifer, MD, FACC, FAHA; Mark S. Link, MD, FACC, FHRS; Srihari S. Naidu, MD, FACC, FSCAI; Rick A. Nishimura, MD, FACC, FAHA; Steve R. Ommen, MD, FACC, FAHA; Harry Rakowski, MD, FACC, FASE; Christine E. Seidman, MD, FAHA; Jeffrey A. Towbin, MD, FACC, FAHA; James E. Udelson, MD, FACC, FASNC; Clyde W. Yancy, MD, FACC, FAHA

American College of Cardiology Foundation/American Heart Association (ACCF/AHA) Task Force Members: Alice K. Jacobs, MD, FACC, FAHA (*Chair, 2009–2011*); Sidney C. Smith, JR, MD, FACC, FAHA (*Immediate Past Chair, 2006–2008*)*; Jeffrey L. Anderson, MD, FACC, FAHA (*Chair-Elect*); Nancy M. Albert, PhD, CCNS, CCRN, FAHA; Christopher E. Buller, MD, FACC*; Mark A. Creager, MD, FACC, FAHA; Steven M. Ettinger, MD, FACC; Robert A. Guyton, MD, FACC; Jonathan L. Halperin, MD, FACC, FAHA; Judith S. Hochman, MD, FACC, FAHA; Harlan M. Krumholz, MD, FACC, FAHA*; Frederick G. Kushner, MD, FACC, FAHA; Rick A. Nishimura, MD, FACC, FAHA*; E. Magnus Ohman, MD, FACC; Richard L. Page, MD, FACC, FAHA*; William G. Stevenson, MD, FACC, FAHA; Lynn G. Tarkington, RN*; Clyde W. Yancy, MD, FACC, FAHA

*Former Task Force member during this writing effort.

Financial Disclosures/Conflicts of Interest

The Task Force makes every effort to avoid actual, potential, or perceived conflicts of interest that may arise as a result of relationships with industry and other entities (RWI) among the writing committee. Specifically, all members of the writing committee, as well as peer reviewers of the document, are required to disclose all relevant relationships and those 12 months prior to initiation of the writing effort. The policies and procedures for RWI for this guideline were those in effect at the initial meeting of this committee (March 28, 2009), which included 50% of the writing committee with no relevant RWI. All guideline recommendations require a confidential vote by the writing committee and must be approved by a consensus of the members voting. Members who were recused from voting are indicated on the title page of this document with detailed information included in Appendix 1 in the original guideline document. Members must recuse themselves from voting on any recommendations where their RWI apply. If a writing committee member develops a new RWI during his/her tenure, he/she is required to notify guideline staff in writing. These statements are reviewed by the Task Force and all members during each conference call and/or meeting of the writing committee and are updated as changes occur. For detailed information regarding guideline policies and procedures, please refer to the ACCF/AHA methodology and policies manual. RWI pertinent to this guideline for authors and peer reviewers are disclosed in Appendixes 1 and 2 in the original guideline document, respectively.

Guideline Status

This is the current release of the guideline.

Guideline Availability

Electronic copies: Available in Portable Document Format (PDF) from the [American College of Cardiology Web site](#) and the [Circulation Web site](#) .

Print copies: Available from the ACC, 2400 N Street NW, Washington DC, 20037; (800) 253-4636 (US only).

Availability of Companion Documents

The following are available:

- Gersh BJ, Maron BJ, et al. 2011 ACCF/AHA guideline for the diagnosis and treatment of hypertrophic cardiomyopathy: executive summary. Bethesda (MD): American College of Cardiology/American Heart Association. 2011 Dec. 38 p. Electronic copies: Available in Portable Document Format (PDF) from the [Journal of the American College of Cardiology \(JACC\) Web site](#) .
- ACCF/AHA pocket guideline. Guideline for the diagnosis and treatment of hypertrophic cardiomyopathy. Bethesda (MD): American College of Cardiology/American Heart Association. 2011 Nov. 42 p. Electronic copies: Available in PDF from the [JACC Web site](#) .
- 2011 ACCF/AHA guideline for the diagnosis and treatment of hypertrophic cardiomyopathy. Slide set. Bethesda (MD): American College of Cardiology/American Heart Association. 2011. 74 p. Electronic copies: Available from the [JACC Web site](#) .
- Methodology manual and policies from the ACCF/AHA Task Force on Practice Guidelines. 2010 Jun. 88 p. American College of Cardiology Foundation and American Heart Association, Inc. Electronic copies: Available in PDF from the [American College of Cardiology \(ACC\) Web site](#) .

Print copies: Available from the ACC, 2400 N Street NW, Washington DC, 20037; (800) 253-4636 (US only).

Patient Resources

None available

NGC Status

This NGC summary was completed by ECRI Institute on April 18, 2012. The information was verified by the guideline developer on May 16, 2012. This summary was updated by ECRI Institute on January 23, 2013 following the U.S. Food and Drug Administration advisory on Pradaxa (dabigatran etexilate mesylate).

Copyright Statement

This NGC summary is based on the original guideline, which is subject to the guideline developer's copyright restrictions as follows:

Copyright to the original guideline is owned by the American College of Cardiology Foundation (ACCF) and the American Heart Association, Inc. (AHA). NGC users are free to download a single copy for personal use. Reproduction without permission of the ACC/AHA guidelines is prohibited. Permissions requests should be directed to copyright_permissions@acc.org

Disclaimer

NGC Disclaimer

The National Guideline Clearinghouse[®],[®] (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at <http://www.guideline.gov/about/inclusion-criteria.aspx>.

NGC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI Institute, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.